FORMULA 273

FFE TERG-O-SAN CLEANER-SANITIZER

HEAVY DUTY FORMULA FOR CLEANING,
DEGREASING AND SAMITIZING MEAT DEPARTMENT, BAKERY, AND DELICATESSEN EQUIPMENT

ACTIVE INGREDIENTS:	
n-Alkyl (60% C_{14} , 30% C_{16} , 5% C_{12} , 5% C_{18})	
Dimethyl Benzyl Ammonium Chlorides	0.75%
n-Alkyl (50% C ₁₂ , 30% C ₁₄ , 17% C ₁₆ , 3% C ₁₈)	
Dimethyl Ethylbenzyl Ammonium Chlorides	0.75%
Trisodium Phosphate	2.00%
Sodium Metasilicate, Anhydrous	2.00%
Trisodium Salt of N-hydroxyethylethylenediamine	
triacetic acid	0.04%
INERT INGREDIENTS (Includes cleaning agents,	
penetrants and emulsifiers)	94.46%
EPA Reg. No. 1685-67 TOTAL 1	00.00%

WARNING KEEP OUT OF REACH OF CHILDREN

See right panel for antidote statement and other precautions.



DIRECTIONS FOR USE

It must be diluted with either hot or cold water before using.

Meat, Bakery, Delicatessen and Fish and Chips equipment or utensils: First remove or protect food products and packaging materials. Dilute four ounces of FFC Terg-O-San Cleaner-Sanitizer per gallon of water for sanitizing action against Staphylococcus aureus and Escherichia coli. Wash equipment with the diluted solution utilizing brush. Follow with a rinse of clear (Potable) water. For heavy grease conditions use stronger solution and hot water for better degreasing action.

Floors, tables, walls and other hard surface cleaning and sanitizing: Dilute four ounces of FFC Terg-O-San Cleaner-Sanitizer per gallon of water and apply with mop or cloth.

Rinse food contact surfaces with terminal potable water before reuse.

FFC Terg-O-San Cleaner-Sanitizer is effective in water up to 400 ppm hardness calculated as CaCo₃.

WARNING KEEP OUT OF REACH OF CHILDREN

Causes severe eye irritation. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food. Rinse empty container with water before discarding.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse. If swallowed drink promptly a large quantity of milk, egg whites, geletin solution or if they are not available drink large quantities of water. Avoid alcohol. Call a physician.

NET CONTENTS

Feb-15-1975

160