

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wash hands after using. Do not contaminate food or feedstuffs with this spray. Do not allow spray to contact food, feedstuffs or water supplies. Thoroughly wash food contact surfaces with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying. Do not use this in edible product areas of food processing plants, restaurants or other food handling areas while food processing is underway. Cover or remove any food or food processing equipment during application.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get immediate medical attention.
IF IN EYES OR ON SKIN: Flush with plenty of water for at least 15 minutes. For eyes, get medical attention.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. If symptoms of cholinesterase inhibition appear, atropine only by injection is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL AND CHEMICAL HAZARDS

Do not apply this water-based product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. Do not use on surfaces that can be harmed or stained by water.

DIRECTIONS FOR USE

It is violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
STORAGE: Do not store at temperatures below 32°F.
PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.
CONTAINER DISPOSAL: Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved State and Local procedures.

For use in nonfood areas indoors and outside premises of hotels, motels, homes, preschools, apartment buildings, commercial office, institutional and storage buildings, recreational buildings, boats, ship cabins, and hospitals. In hospitals, remove patients prior to treatment. Ventilate room for two hours before returning patients.

This product can be used in nonfood areas only, such as corridors, offices, foyers, lavatories, garages, utility rooms, storage rooms and basements of packaging and processing plants, milk handling areas of milk plants, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, and bakeries.

PH 12 1685-4

State FORMULA 401
**READY KILL
with DURSBAN***

A combination of insecticides offering good flushing activity, quick knockdown and residual kill of cockroaches for up to 4 weeks

For residual and contact control of cockroaches (waterbugs), crickets, silverfish, clover mites, centipedes, millipedes, firebrats, earwigs, sowbugs, spiders, brown dog ticks, lone star ticks, ants, fire ants, carpet beetles, fleas, boxelder bugs, scorpions, bedbugs, cigarette beetles, chocolate moths and clothes moths

STIR OR SHAKE WELL BEFORE USING

ACTIVE INGREDIENTS:

*Resmethrin [5-(phenylmethyl)-3-furanyl] methyl-2,2-dimethyl-3-(2-methyl-1-propenyl) cyclopropanecarboxylate 0.10%

*Chlorpyrifos 0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate 0.50%

INERT INGREDIENTS 99.40%

*Cis/trans isomers ratio: max 30% (±) cis and min 70% (±) trans

†U.S. Patent Nos. 3,465,007 and 3,542,928 and 3,683,078

*Penick Corp's SBP-1382® brand of Resmethrin insecticide

*U.S. Patent No. 3,244,585. Dursban is a registered trademark of Dow Chemical Company.

PRECAUCION AL USUARIO: Si usted no lee ingles, no use esta product haste que la etiqueta le haya sido explicada ampliamente.
TRANSLATION (To The User: If you cannot read English, do not use this product until the label has been fully explained to you).

KEEP OUT OF REACH OF CHILDREN

CAUTION

**SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

IN FOOD HANDLING ESTABLISHMENTS, FORMULA 401 READY KILL WITH DURSBAN* CAN BE USED IN NON-FOOD AREAS AS A CONTACT OR RESIDUAL SPRAY. IN FOOD AREAS, APPLICATION IS LIMITED TO SPOT AND OR CRACK AND CREVICE TREATMENT.

The **State** CHEMICAL MANUFACTURING CO.
MILWAUKEE, WISCONSIN

EPA REG NO 1685-44

EPA Est No 1685-GH-1
NET CONTENTS

48 For Industrial Use Only
ACCEPTED
APR 12 1984
U.S. Environmental Protection Agency
EPA REG NO 1685-44

Also, for use in and around campers, mobile homes, buses and cars. This product can be used in cargo planes, trucks, boxcars and ship cargo holds only if the vehicles are not used for transport of food or food products at any time.

Also, for use in dog kennels and horse stables. Do not spray on horses and dogs. Do not spray on dog and horse food or feed items.

One gallon of spray is sufficient to treat 1500 linear feet with a 8 inch spray. For contact and residual spray, use State Formula 401 Ready Kill with Dursban* without diluting.

CONTACT SPRAY KILLS AND REDUCES INFESTATIONS: Use as a spot treatment to kill the insects listed on the center panel of this label. When possible spray directly on pests to reduce infestations. Apply as a coarse spray or with a paint brush into hiding places such as cracks, crevices, moist area around pipes and sinks, under refrigerators and other appliances, baseboards and storage areas. Apply spray until area is wet, but not to the point of runoff. Repeat as necessary.

FLEAS, BROWN DOG AND LONE STAR TICKS: Thoroughly apply as a spot treatment to infested areas such as pets' beds and resting quarters, nearby cracks and crevices, along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment of pet areas. **DO NOT TREAT PETS WITH THIS PRODUCT.** To control the source of your flea infestation, treat your pets with a product registered for animal application, such as State 797-A Powdered Insecticide.

CARPET BEETLES AND CLOTHES MOTHS: Do not spray clothing with this product. Do not allow clothing to contact areas where this product has been used. Spray edges of carpeting and under carpeting, rugs and closets. Make localized application to the floor and baseboards. Spray directly into cracks, crevices, closets and infested areas of shelving and/or wherever these insects are seen or suspected. Repeat as necessary.

RESIDUAL SPRAY: Aids in the control of cockroaches, crickets, silverfish, centipedes, millipedes, earwigs, sowbugs. Kills up to 4 weeks. Apply as for contact spray. In basements, apply in hiding places, in cracks and crevices, lower walls and dark places. Spray surfaces until wet, but not to the point of run-off. Treat on a monthly basis or as necessary. **OUTDOOR PESTS**

PESTS ON OUTSIDE SURFACES OF BUILDINGS: Use State Formula 401 Ready Kill with Dursban* to control ants, clover mites, crickets, earwigs, millipedes, and sowbugs by application as a residual spray or with a paint brush outside surfaces of buildings including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Do not treat or allow the spray to contact plants (lawns, flowering plants, shrubs, trees, evergreens, etc.) because of possible injury. Repeat treatment as needed to maintain effectiveness.

NOTE: To help prevent infestation of buildings by ants, crickets, earwigs, and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the buildings, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance.

TO CONTROL FIRE ANTS: Thoroughly wet the mound and surrounding area to a 4 foot diameter with Formula 401 Ready Kill with Dursban*. For best results apply in cool weather, 65-80°F or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity and treat all colonies which may not as yet have constructed a mound.