

6075-555-4712

RODENTICIDE, BAIT, ANTICOAGULANT
FED. SPEC. C-R-00000 (INT-FWS)
24 Ounces

ACTIVE INGREDIENTS: 2 pivalyl 1,3-indandione	0.005%
INERT INGREDIENTS	99.995%
Total	100.000%

DIRECTIONS:

Place 4 to 8 ounces of Rodenticide, Bait, Anticoagulant in bait stations. Place so as to protect children, pets, and livestock in feeding areas where rats or mice feed. Refresh baits as necessary, and maintain bait stations for not less than 12 days. (Baiting may be continued indefinitely, and increased time of exposure of baits may result in better control of rodents.) If bait molds, decomposes, or becomes sour, replace with fresh material.

CAUTION:

Keep away from humans, domestic animals, and pets. If swallowed by humans, domestic animals, or pets, this material may reduce the coagulability of the blood and cause bleeding. In case of poisoning, induce vomiting and administration of Vitamin K₁ is indicated. In case of hemorrhage, transfusion is indicated as in the case of hemorrhage caused by overdosage of dicoumatil, diphacinone.

U.S. DEPT. OF AGRICULTURE, OFFICE OF PEST CONTROL ADMINISTRATION

CHEMICAL COMPANY, INC.
JERSEY CITY, N.J.

USDA Reg. No. 1691-31