

RODENTICIDE, BAIT
ANTICOAGULANT PMF

USDA Reg. No. 1691-88

For Control of Commensal Rats and Mice

Stock No. 6840-753-4973

ACTIVE INGREDIENTS:

2-Isovaleryl-1,3-Indandione .0004

INERT INGREDIENTS:

Liquid Petrolatum	4.8	%	
Sucrose	4.8	%	
Dye (G#305 Flaming Red-OX Color Works)	0.05	%	
Breakfast Rolled Oats	90.200	%	

DIRECTIONS: Place 4 to 8 ounces of Rodenticide, Bait, Anticoagulant in bait stations. Place so as to protect children, pets, and livestock in locations where rats or mice feed. Maintain bait stations uninterruptedly for not less than 15 days, replenishing as necessary. Baiting may be continued indefinitely where re-infestation by rats and mice is a problem. If bait molds, decomposes, or becomes sour, replace with fresh material.

For additional details see tri-service manual, Insect and Rodent Control, AFM 75-7, TMS-632, NAVDOCKS 11-Pc-2.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Keep away from humans, domestic animals, and pets. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases intravenous and oral administration of Vitamin K combined with blood transfusion are indicated as in the case of hemorrhage caused by overdose of dihydroxyacetone.

Manufactured by
CHEMICAL COMPOUNDING CORPORATION
532 Johnston Avenue
Jersey City 4, N.J.

FIVE POUNDS NET