

SANTROL

Sanitizing Agent

No. 25

Active Ingredients:
n-Alkyl (C₁₄-50%, C₁₂-40%, C₁₆-10%)
dimethyl benzyl ammonium chlorides...10%
Inert Ingredients.....90%

Phenol Coefficient of 50 for S Typhosa at
20 degrees C.
EPA Reg. No. 1696-22

DANGER

**KEEP OUT OF REACH OF
CHILDREN**

HARMFUL IF SWALLOWED

Do not get in eyes. For eye contact, flush thoroughly and immediately get medical attention. Avoid contact with skin. In case of contact, flush with plenty of water. If irritation persists, get medical attention. Avoid contamination of food. Rinse empty container thoroughly with water and discard it.

METRO PRODUCTS CO.

ATLANTA, GA. 30336
NET CONTENTS 1 GAL.

DIRECTIONS: Thoroughly clean all objects and surfaces before disinfecting with SANTROL solution. **IMPORTANT:** soaps and anionic detergents reduce the germicidal efficiency of SANTROL so care should be taken to rinse objects thoroughly before placing in SANTROL solution. Non-ionic detergents do not affect the efficiency of SANTROL.

FOR GENERAL DISINFECTION OF POULTRY HOUSES, COOPS, ETC.:

1. Remove all poultry and feeds from premises, trucks, coops, or crates.
2. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry.
3. Empty all troughs, racks and other feeding and watering appliances.

Continued

4. Saturate surfaces with a solution containing 1 ounce of SANTROL per 2 gallons of water for non-porous surfaces or 2 ounces of SANTROL per 2 gallons of water for porous surfaces. This may be done by spraying, mopping, or wiping with sponges or rags.

5. Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried.

6. All treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be thoroughly rinsed with potable water prior to reuse.

FOR DISINFECTION OF HATCHING TRAYS:

Immerse previously cleaned trays in a solution containing 1 ounce of SANTROL to each 2 gallons of water.

