

KLEEN-SAN HATCHERY CLEANER NO. 45

ACTIVE INGREDIENTS:

Alkyl (C ₁₁ 50%, C ₁₂ 40%, C ₁₆ 10%) Dimethyl benzyl ammonium chlorides	5.0%
Sodium Carbonate	2.75%
Sodium Metasilicate	1.29%
Trisodium Phosphate	.97%
Tetrasodium ethylene diamine tetraacetate	.39%
INERT INGREDIENTS	89.6%

Note: Contains Sodium Nitrite

DIRECTIONS

FOR CLEANING AND SANITIZING HATCHERS: Dilute 1 Oz. Kleen-San Hatchery Cleaner No. 45 to each gallon of water. Using this solution and a stiff brush, vigorously scrub the surface to be cleaned. Then sponge clean with clear water. Rinse bucket and brush, and use a fresh solution of Kleen-San Hatchery Cleaner No. 45 for each hatcher.

FOR CLEANING AND SANITIZING HATCHING TRAYS: Soak trays in a solution containing 1 Oz. of Kleen-San Hatchery Cleaner No. 45 to each gallon of water for 2 to 5 minutes. Then vigorously brush clean and rinse in clear water.

FOR CLEANING AND SANITIZING FLOORS: Use 1 cup Kleen-San Hatchery Cleaner No. 45 to 3-4 gallons water. Spread liberally but do not flood. Pick up with a mop, squeegee, or vacuum cleaner.

FOR DISINFECTION: Use a solution containing 1 Oz. Kleen-San Hatchery Cleaner No. 45 to each gallon of water as a final rinse after surfaces have been thoroughly cleaned.

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food. Dilute with water as directed. Do not mix with soap or anionic detergent. Rinse empty container thoroughly and discard it.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink **promptly** a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

E. P. A. Reg. No. 1696-11

Manufactured by
METRO PRODUCTS COMPANY
Manufacturing Chemists
Atlanta, Georgia

Aug 11-1977

Net Contents _____ Gallons

1696-11