

DIRECTIONS FOR USE

EATING ESTABLISHMENTS, RESTAURANTS, CAFES and TAVERNS: (1) Wash utensils with a good detergent\*, (2) rinse with clear water, (3) dip in SANTROL solution 200 ppm for 2 minutes or for the contact time specified by the governing sanitary code.

MILK FARMS and MILK PROCESSING PLANTS: First clean equipment and utensils thoroughly with a good cleansing agent\*. Then sanitize with SANTROL 200 ppm. A further rinse in clean running water is recommended.

SANTROL is not to be used as a preservative for foods.

FOOD PROCESSING PLANTS: BAKERIES, CANNERIES, etc.: Clean walls and floors with a cleansing agent\*; then spray with S A N T R O L 800 ppm. Clean equipment thoroughly with a cleansing agent\* then sanitize by wiping or spraying SANTROL 200 ppm. Before direct contact with food, a rinse with clean running water is necessary.

\*When soap or an anionic detergent is used for cleaning, rinse with water before sanitizing.

# SANTROL

## Sanitizing Agent

Active Ingredients:

n-Alkyl (C <sub>11</sub> 50%, C <sub>12</sub> 40%, C <sub>16</sub> 10%) dimethyl benzyl ammonium chlorides	10.0%
Ethyl Alcohol	2.5%
Inert Ingredients	87.5%
Phenol Coefficient of 50 for S. Typhosa at 20° C	

### DANGER:

KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food. See first aid and other precautions on right panel. Rinse empty container thoroughly with water and discard it.

EPA Registration No. 1696-46

Net Contents 1 Gal.

**METRO PRODUCTS COMPANY**  
4720 FREDERICK DR., S. W., ATLANTA, GEORGIA 30336  
EPA Est. No. 1696-GA1

HOSPITALS AND HOUSEHOLD USE: For the general disinfection of floors, walls, furniture, bath tubs, stall showers, toilet seats, etc., first clean with a good detergent\*. Rinse or spray cleaned surfaces with SANTROL 800 ppm.

DIRECTIONS FOR DILUTION FOR SANTROL 200 ppm. approx. add and stir one ounce (2 tablespoonfuls) of SANTROL 10% in 4 gallons of water.

FOR SANTROL 800 ppm. approx. add and stir one ounce (2 tablespoonfuls) of SANTROL 10% in one gallon of water.

Sanrol is effective in water up to 550 ppm hardness calculated as CaCO<sub>3</sub> by the AOAC Germicidal Detergent Sanitizer Test.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

