

WINRU

75% EMULSIFIABLE CHLORDANE CONCENTRATE

A water dispersable chlordane concentrate containing 75% technical chlordane.

ACTIVE INGREDIENTS:

*Technical Chlordane	. 75%
Petroleum Distillate	10.00
INERT INGREDIENTS:	15%
*Equivalent to 45% of Octachloro-4.7-Methanotetrahydroindane	1002

and 30% of related compounds.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Avoid inhalation and skin contract. In case of spillage on skin wash with soap and water. Avoid contamination to feed and foodstuffs.

This solution if brought into contact with the skin can cause toxic symptoms.

This insecticide should not be stored, sprayed or otherwise exposed toward or near an open flame.

See right panel for additional cautions.

D'RECTIONS:

Subterranean Termites: Winru 75% Emulsifiable Chlordane may be used as an aid in the control of Subterranean termites.

Ground Treatments: Dilute Winru 75% Emulsifiable Chlordane at the rate of 1 part concentrate to 49 parts water and apply in trench at the rate of one gallon of mixed spray per linear foot. Trench should be 30" deep around foundations, but not below footings. Be sure to treat entire area around buildings. Annual inspection should be made for possible reinfestations. Winru 75% Emulsifiable Chlordane when diluted as directed above will not cause phytoxicity.


Roaches: To control exposed stages apply as a 2% emulsion (14 oz. in 5 gallons of water) spraying thoroughly in all cracks and crevices and other hiding places. Repeat treatment as needed.

Ants: Apply as a 2% emulsion directly to ant trails and areas around door sills and window frames.



GALLONS

Manufactured by WINRU CHEMICAL & SALES COMPANY CHEMICAL & SALES COMPANY 64101



DIRECTIONS (Continued):


Buildings. Wirru 75% Emulsifiable Ch! applied is a residual spray in barns, stan and similar buildings. It is effective ants, cockroaches, brown dog ticks, all particular attention to cracks in stalls an quarters of animals. Apply as a 2% so 5 gallons of water) making application weeks or oftener if exposed to weatheri. Do not use in dairy barns, milk houses, grain or bulk food storage bins.


CAUTION:

Use only according to label directions NOT FOR HOUSEHOLD USE. DO NOT FRODUCTS AREAS OF FOOD PROCES: Do not contaminate water by cleaning disposal of wastes.

Do not reuse empty drum. Return to dr or destroy by perforating or crushing a safe place.

Serving the FOOD INDUSTRY





WINRU

75% EMULSIFIABLE CHLORDANE CONCENTRATE

fiable Chlorol of Subter-

Emulsifiable e to 49 parts one gallon of be 30" deep s. Be sure to al inspection s. Winru 75% irected above

as a 2% emulng thoroughly niding places.

to ant trails frames.

*Technical Chlordane 75° Petroleum Distillate 10° INERT INGREDIENTS: 15° *Equivalent to 45% of Octachloro-4,7-Methanotetrahydroindane 100°	A water dispersable chlordane concentrate containing 75% technic ACTIVE INGREDIENTS:	cal chlordane
Petroleum Distillate 10° INERT INGREDIENTS: 15° *Equivalent to 45% of Octachloro-4,7-Methanotetrahydroindane 100°		75 [°]
*Equivalent to 45% of Octachloro-4,7-Methanotetrahydroindane 100°		
	INERT INGREDIENTS:	. 15%
and do 76 of tetated composites.	*Equivalent to 45% of Octachloro-4,7-Methanotetrahydroindane and 30% of related compounds.	100,

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Avoid inhalation and skin contract. In case of spillage on skin wash with soap and water. Avoid contamination to feed and foodstuffs.

This solution if brought into contact with the skin can cause toxic symptoms.

This insecticide should not be stored, sprayed or otherwise exposed toward or near an open flame.

See right panel for additional cautions.

NET CONTENTS

GALLONS

Manufactured by MANURU CHEMICAL & SALES COMPANY MINES LINE, KANSAS CITY, MISSOURS 64101



DIRECTIONS (Continued):

Buildings. Wirry 75% Emulsifiable Chlordane may be applied is a residual spray in barns, stables, dog houses, and similar buildings. It is effective for controlling ants, cockroaches, brown dog ticks, and spiders. Pay particular attention to cracks in stalls and other sleeping quarters of animals. Apply as a 2% solution (14 oz. in 5 gallons of water) making applications every 4 to 6 weeks or oftener if exposed to weathering.


Do not use in dairy barns, milk houses, poultry houses, grain or bulk food storage bins.

CAUTION:

Use only according to label directions and limitations. NOT FOR HOUSEHOLD USE. DO NOT USE IN EDIBLE FRODUCTS AREAS OF FOOD PROCESSING PLANTS.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.



TRANSFERRED FROM