

DIRECTIONS

DIRECTIONS FOR USE OF WINRU WATER SOLUBLE WARFARIN: One Third ounce of Winru Water Soluble Warfarin will make one quart of solution. Drinking water in a Mason jar can be used. No mixing or stirring required. **DO NOT ATTEMPT TO DISSOLVE SMALL SILICA PARTICLES** which remain on the bottom of the jar. Place a chicken water fount over the mouth of the jar and turn over quickly. ~~Set fount on a clean level surface where rats are accustomed to feed. Discard at once water in which mold or filth is noted. Keep fount clean and well supplied at all times. Do not place fount where freezing may occur. Protect founts from children or pets and domestic animals.~~

Where a continuous source of infestation prevails from nearby dumps or fields, permanent bait stations should be used and baits replenished as needed.

MICE: For controlling mice follow the same general directions as for rats except bait placements may be smaller and more placements should be made. Mice are more difficult to control than rats and complete control may take a longer period of baiting.

IMPORTANT— BAITING SHOULD CONTINUE UNTIL COMPLETE LACK OF FEEDING IS NOTED. THIS SHOULD BE FROM 5 TO 14 DAYS. It is recommended that Warfarin Prepared Baits be used in conjunction with this product, especially where there is an ample supply of fresh water.

CAUTION: Winru Water Soluble Warfarin contains as its active ingredient an anti-coagulant chemical which if taken ~~accidentally~~ by humans, domestic animals, or pets may reduce the clotting ability of the blood and serious hemorrhage may result. In case baits are accidentally swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

NOTE FOR PHYSICIANS: When a human has been known to have accidentally ingested Winru Water Soluble Warfarin, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol

WINRU Water Soluble WARFARIN

A concentrate of Sodium Salt of Warfarin*

FOR USE AGAINST RATS AND MICE

Active Ingredients:

Sodium Salt of Warfarin (Sodium Salt of 3-(α -acetonylbenzyl)—4-hydroxycoumarin) 0.54%**

Inert Ingredients:

(Silica) 99.46%
Total 100.00%

*Wisconsin Alumni Research Foundation—patent applied for.
**Equivalent 0.5% Warfarin.

Distributed By

Winru Chemical & Sales Co.

923 STATE LINE
KANSAS CITY, MO.

NET CONTENTS: 1 3 OUNCE

CAUTION: Keep out of reach of children. Read caution and antidote statement on back panel.

U.S. PAT. NO. 1700-E

4002
UNDEVELOPED
FOR RECORDS
FD-302 (REV. 1-25-60)
7700-174 SUBJECT
TO ATTACHED COMMENTS.

WINRU WATER SOLUBLE WARFARIN

Corrected Directions For Use

1. Let point on clean, level surface where rats are accustomed to feed. Discard at once water in which hold or place bait. Keep point of baiting clean and dry at all times. Baiting should be done in frequent way occur. Baiting should be continued for several days until all rats are eliminated.

2. Baiting should be done in distribution of baits from nearby houses or fields, particularly bait should be used. In all cases of ~~XXX~~ infestations baiting should be continued until all rats are eliminated ten days.

3. Baiting should be done in all cases where general directions as for rats are not sufficient. Baiting should be done in all cases of infestations of houses. Baiting should be done in all cases of infestations of houses. Baiting should be done in all cases of infestations of houses.

4. Baiting should be done in all cases of infestations of houses. Baiting should be done in all cases of infestations of houses. Baiting should be done in all cases of infestations of houses.

5. Baiting should be done in all cases of infestations of houses. Baiting should be done in all cases of infestations of houses. Baiting should be done in all cases of infestations of houses. Baiting should be done in all cases of infestations of houses.

DIRECTIONS

DIRECTIONS FOR USE OF WINRU WATER SOLUBLE WARFARIN: One-third ounce of Winru Water Soluble Warfarin will make one quart of solution. Drinking water in a Mason jar can be used. No mixing or stirring required. **DO NOT ATTEMPT TO DISSOLVE SMALL SILICA PARTICLES** which remain on the bottom of the jar. Place a chicken water fount over the mouth of the jar and turn over quickly. ~~Set fount on a clean level surface where rats are accustomed to feed. Discard at once water in which mold or dirt is noted. Keep fount clean and well supplied at all times. Do not place fount where freezing may occur. Protect founts from children or pets and domestic animals.~~

~~Where a continuous source of infestation prevails from nearby dumps or fields, permanent bait stations should be used and baits replenished as needed.~~

~~MICE: For controlling mice follow the same general directions as for rats except bait placements may be smaller and more placements should be made. Mice are more difficult to control than rats and complete control may take a longer period of baiting.~~

~~**IMPORTANT: BAITING SHOULD CONTINUE UNTIL COMPLETE LACK OF FEEDING IS NOTED. THIS SHOULD BE FROM 5 TO 14 DAYS.** It is recommended that Warfarin Prepared Baits be used in conjunction with this product, especially where there is an ample supply of fresh water.~~

CAUTION: Winru Water Soluble Warfarin contains as its active ingredient an anti-coagulant chemical which if taken ~~continually~~ by humans, domestic animals, or pets may reduce the clotting ability of the blood and serious hemorrhage may result. In case baits are accidentally swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

NOTE FOR PHYSICIANS: When a human has been known to have accidentally ingested Winru Water Soluble Warfarin, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol.

WINRU Water Soluble WARFARIN

A concentrate of Sodium Salt of Warfarin*

FOR USE AGAINST RATS AND MICE

Active Ingredients:

Sodium Salt of Warfarin (Sodium Salt of 3-(α -acetylbenzyl)-4-hydroxycoumarin) 0.54%**

Inert Ingredients:

(Silica) 99.46%
Total 100.00%


*Wisconsin Alumni Research Foundation—patent applied for.
**Equivalent 0.5% Warfarin.

Distributed By
Winru Chemical & Sales Co.
923 STATE LINE
KANSAS CITY, MO.

NET CONTENTS: 1/3 OUNCE

CAUTION: Keep out of reach of children. Read caution and antidote statement on back panel.

USDA Reg. No. 1700-E



14 PT. TYPE

(SEE ATTACHED CORRECTED DIRECTIONS)