09/30/1964

CUPROSE (R)

## Algaecide

## ACTIVE INGREDIENTS:

\* Inert ingredients contain stabilizer

Copper expressed as metallic....20.4%

DIRECTIONS: For control of algae in industrial ponds feed 2 to 15 ppm Cuprose; for reservoirs do not exceed 13.3 lbs./acre foot (equivalent to 1 ppm as Cu) applied by broadcasting or use of mechanical spreader. ease note that 1 ppm copper may kill some fish.

For control of algae in industrial cooling water systems use 1 to 5 ppm Cuprose (0.2 to 1.0 ppm copper content as Cu). Feed Cuprose either as a 5% water solution through plastic or ceramic proportioning equipment or dry through stoker type feeder. Application may be continuous or intermittent depending upon system characteristics. Further directions for use obtainable from Nalco.

NOTE: Maximum copper content permissible in finished potable water supplies is 1 ppm according to 1962 Public Health Service Drinking Water Standards.

CAUTION: HARMFUL IF SWALLOWED. Avoid contact with eyes. Avoid prolonged or repeated contact with skin. Avoid breathing dust. In case of contact, wash skin with soap and water; for eyes, immediately 'ush with large amounts of water for at least 15 minutes, and get medical attention.

## KEEP OUT OF REACH OF CHILDREN

Seller makes no warranty of any kind, express or implied, concerning the use of this product. Buyer or user assumes all risk of use or handling, whether in accordance with directions or not.

## NALCO CHEMICAL COMPANY

6216 West 66th Place

Chicago, Illinois 60638

Net Contents Shown Elsewhere On Container

Form No. 4.115.002

ACCEPTED

SEP 30 1964

UNDER THE FEDERAL INSECTICIDE FUNCTCIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER.