



NALCO[®]

321

ACCEPTED

OCT 9 1975

UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND ANTIMOLD ACT
FOR ECONOMIC POISON REGISTERED
EPA UNDER NO. 1706-140

MICROORGANISM CONTROL CHEMICAL

ACTIVE INGREDIENT:

1-Alkyl (C₆ to C₁₈)* amino-3-aminopropane diacetate 23.5%

INERT INGREDIENTS: 76.5%

Total 100.0%

* As in fatty acid of coconut oil.

Weight per gallon = 7.85 lb
EPA Registration No. 1706-140

NALCO CHEMICAL COMPANY

6216 WEST 66TH PLACE CHICAGO, ILLINOIS 60638

EPA est: 1706-IL-1

NET CONTENTS SHOWN ELSEWHERE ON CONTAINER

DIRECTIONS: For bacterial and fungal slime control in industrial recirculating cooling water systems use 150—200 ppm (1.5 to 2.0 gallons/10,000 gallons of system water) NALCO 321 initially. Repeat as needed to achieve control. Badly fouled systems must be precleaned. Once the desired degree of control has been achieved apply 50—100 ppm (0.5 to 1.0 gallons/10,000 gallons of system water) NALCO 321 weekly or as needed to maintain control.

Feed NALCO 321 directly from the drum to a point in the system where it will be evenly mixed and uniformly distributed. Foaming may occur at the higher end of the dosage range. To control foaming use 5—100 ppm NALCO 71-D5 antifoam or other suitable antifoam.

NOTE: NALCO 321 INTENDED FOR INDUSTRIAL USE ONLY. Do not apply to potable water or domestic water systems.

This product is toxic to fish. Do not discharge treated effluent where it will drain into lakes, streams, ponds or public water.

Do not reuse this container. Destroy when empty. Apply this product only as specified on this label.

DANGER: CORROSIVE. KEEP OUT OF THE REACH OF CHILDREN. CAUSES EYE AND SKIN DAMAGE. DO NOT USE. POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Keep container closed when not in use. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. May be harmful if swallowed, inhaled or absorbed through skin. Use with adequate ventilation. Do not take internally.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg white, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician at once.

NOTE TO PHYSICIAN: Probably mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsions may be needed.