

- **DEEP CLEANS**
- **DEODORIZES**
- **DISINFECTS**

Phenol Coeff: S. aureus—32
S. typhosa—42

**CONCENTRATED
DISINFECTING CLEANER
& DEODORIZER**

HY-GARD

ACTIVE INGREDIENTS: n-Alkyl (50% C14, 40% C12, 10% C16) dimethyl benzyl ammonium chloride 5.0%, Sodium metasilicate 3.0%, Tetrasodium salt of ethylene diamine tetraacetic acid 1.8%. Essential oils 0.30%.

INERT INGREDIENTS: 89.90%. Includes detergents, certified colors and other grease cutting agents.

DANGER: KEEP OUT OF REACH OF CHILDREN

E P A Reg. No. 1712-6

See First Aid Statement and other precautions on back panel.

HYDROL CHEMICAL CO., PHILA., PA. 19104—SINCE 1929

**DISINFECTANT—CLEANER—SANITIZER
USE DIRECTIONS**

FOR CLEANING

Lightly soiled areas—surfaces to be cleaned mopped or brushed with a solution containing 1 ounce of HY-GARD per gallon of water. Rinse cleaned surface with fresh water.

Heavily soiled areas—mop or brush soiled surface with a solution containing 2-3 ounces of HY-GARD per gallon of water. Rinse cleaned surface with fresh water.

GENERAL PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces in homes, schools, restaurants, rest rooms and public areas, mop or brush surface to be treated with a solution containing 1 ounce of HY-GARD per gallon of water. Surfaces that are heavily soiled should be cleaned with 1-2 ounces of HY-GARD per gallon of water and then disinfected. Those surfaces which later come in contact with food must be rinsed with clear water of drinking quality.

To disinfect locker rooms, shower stalls, restrooms, around pools and other areas against growth of fungi, mop area with a solution containing 1/2 ounce of HY-GARD per gallon of water. This solution will destroy *Staphylococcus aureus*, a causative agent of athlete's foot.

HOSPITAL DISINFECTION

To disinfect walls, floors, furniture and other surfaces in hospitals and nursing homes, mop or brush surface to be treated with a solution containing 2 ounces of HY-GARD per gallon of water. Those surfaces that are heavily soiled with blood, fecal matter or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water, then rinsed with a fresh solution containing 2 ounces of HY-GARD per gallon of water before disinfection.

SANITIZATION

To sanitize walls, floors and equipment in processing plants, first clean thoroughly with a solution containing 1/2 ounce of HY-GARD per gallon of water. If surface is heavily soiled, use HY-GARD at 2-3 ounces per gallon to insure thorough cleaning. Sanitize cleaned surface with a rinse solution containing 1/2 ounce of HY-GARD per gallon of water. Food contact surfaces must be rinsed with clear water prior to reuse.

1712-6

DISINFECTANT—CLEANER—SANITIZER—FUNGICIDE—DEODORANT USE DIRECTIONS

Phenol Coeff: *S. aureus*—32
S. typhosa—42

CONCENTRATED CLEANING CLEANER & DEODORIZER HY-GARD®

Alkyl (50% C14, 40% C12, 10% C10), Ammonium chloride 5.0%, Sodium perborate 1.8%, Essential oils 0.30%, Surfactants 90%. Includes detergents, grease cutting agents.

FOR CLEANING

Lightly soiled areas—surfaces to be cleaned should be mopped or brushed with a solution containing 1 ounce of HY-GARD per gallon of water. Rinse cleaned area with fresh water.

Heavily soiled areas—mop or brush soiled surface with a solution containing 2-3 ounces of HY-GARD per gallon of water. Rinse cleaned surface with fresh water.

GENERAL PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces in homes, schools, restaurants, rest rooms and other public areas, mop or brush surface to be treated with a solution containing 1 ounce of HY-GARD per gallon of water. Surfaces that are heavily soiled should be cleaned with 1-2 ounces of HY-GARD per gallon of water and then disinfected. Those surfaces which may later come in contact with food must be rinsed with clear water of drinking quality.

To disinfect locker rooms, shower stalls, walkways around pools and other areas against pathogenic fungi, mop area with a solution containing ¾ ounce in 1 gallon of water. This solution will destroy *Trichophyton interdigitale*, a causative agent of athlete's foot.

HOSPITAL DISINFECTION

To disinfect walls, floors, furniture and other hard surfaces in hospitals and nursing homes, mop or brush surface to be treated with a solution containing 2 ounces of HY-GARD per gallon of water. Those surfaces that are heavily soiled with blood, fecal material or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water and then rinsed with a fresh solution containing 2 ounces per gallon of water before disinfection.

SANITIZATION

To sanitize walls, floors and equipment in food processing plants, first clean thoroughly with a solution containing ¾ ounce of HY-GARD per gallon of water. If surface is heavily soiled, use HY-GARD at a rate of 2-3 ounces per gallon to insure thorough cleaning. Sanitize cleaned surface with a rinse solution containing ½ ounce of HY-GARD per gallon of water. Treated food contact surfaces must be rinsed with potable water prior to reuse.

DEODORIZATION

To deodorize surfaces during cleaning, mop or brush surfaces with a solution containing 2 ounces of HY-GARD per gallon of water. HY-GARD is an effective antibacterial cleaner designed for use in household, industrial and institutional cleaning. When used as directed, HY-GARD will disinfect hard surfaces, such as floors, walls, counter tops and furniture. It may be relied upon to deodorize bathroom areas, garbage cans, food storage areas, kennel areas and other areas where obnoxious odors may develop. This product may be used as a general purpose antibacterial-antifungal detergent in homes, schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other places where efficient cleaning and antimicrobial action is desired. See left panel for use directions.

DANGER

KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eye contact, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of water, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against respiratory depression and convulsions should be taken.

Container Disposal: When empty, rinse container thoroughly with water and destroy it by crushing or burying in a safe place.

E P A Reg. No. 1712-6

anel.

SINCE 1929