

[1737-8]

DIRECTIONS FOR USE

(FOR BEEF CATTLE)

USE IN BACKRUBBERS: To be used undiluted. Use in cable-type backrubbers to protect beef animals from horn flies, mosquitoes and gnats. Fly control with back rubbers is directly proportional to the coverage obtained. The construction and location of the backrubber determines effectiveness. It will receive adequate coverage with the insecticide formulation. The backrubber should be constructed with one end anchored close to ground level and the other end at a height of approximately five feet. The cattle can then treat the underline, shoulders and legs. Locate the backrubber where the animals usually congregate such as near a water trough or salt lick so that it will be used frequently.

CATTLE LICE: If an adequate cable-type backrubber is used throughout the entire year, cattle will not become lousy enough to need spraying. It will usually be necessary to relocate the backrubber to places where cattle loaf during the winter months for best results.

Do not use on dairy cattle or calves under 6 months of age. Do not permit beef animals access to treatment within 28 days of slaughter.

NET CONTENTS ----- GALLON(s)

ACCEPTED
Apr. 3, 1968
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 100-100-000000000000
 SUBJECT TO ATTACHED CONDITIONS

505



CAUTION

KEEP OUT OF REACH OF CHILDREN

See right panel for additional precautions.

READY TO USE.

Aids in controlling horn flies, lice, and gnats on beef cattle.

ACTIVE INGREDIENTS:

Toxaphene*	5.00%
Petroleum oil	95.00%
Total	100.00%

*Technical Chlorinated Camphene Containing 67% to 69% Chlorine
DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

U.S.D.A. Registration No. _____

Sel
or
the
of
wh
tre

303



Additional Precautions

Harmful if swallowed! Avoid prolonged breathing of spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Avoid storage near feed and food products. To protect fish and wildlife, do not contaminate streams, lakes, or ponds with this material.

ANTIDOTE — Internal—Give sodium chloride 15 G (One tablesp on of table salt) in glass of warm water and repeat until vomit is clear. **CALL A DOCTOR IMMEDIATELY.**

External—Wash with soap and water.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

CAUTION

KEEP OUT OF REACH OF CHILDREN

See right panel for additional precautions.

READY TO USE

Aids in controlling horn flies, lice, and gnats on beef cattle.

ACTIVE INGREDIENTS:	per cent by weight
Toxaphene*	5.00%
Petroleum oil	95.00%
Total	100.00%

*Technical Chlorinated Camphene Containing 67% to 69% Chlorine
DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

U.S.D.A. Registration No. _____

Manufactured for and Distributed by

SOUTH OMAHA SUPPLY

(Corporate Name: South Omaha Feed and Supply)

Feed and Farm Supplies

OMAHA, NEBRASKA