

CERTO X

PIVALYL CONCENTRATE

ANTI-COAGULANT RODENTICIDE

CAUTION:

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

2-Pivalyl-1, 3-Indandione50%

INERT INGREDIENTS: 99.50%

TOTAL 100.00%

U.S. PAT. REG. NO. 1749,000

DIRECTIONS: Thoroughly mix one part by weight of PIVALYL CONCENTRATE with 19 parts by weight of a suitable bait material such as oat meal, corn meal, cracked corn, etc.

For increased palatability and to facilitate the preparation of baits, corn, peanut or mineral oil up to 5% and sugar up to 5% may be substituted for equivalent amounts of the bait material.

For top efficiency baits should be placed in areas where rats and mice have been observed feeding. Baits should be placed in locations inaccessible to cats, dogs, livestock and poultry.

GENERAL DIRECTIONS

- a) Place baits in locations where rats and mice feed, water, or travel.
- b) Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable.
- c) Contaminated or spoiled baits should be replaced promptly.
- d) Baiting should continue until all signs of feeding have stopped.

- b) Bait should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes.
- c) An uninterrupted supply of bait should be maintained for at least 15 days.

DIRECTIONS FOR RAT CONTROL

- a) Four ounces of bait per station should be the MINIMUM amount used.

DIRECTIONS FOR MOUSE CONTROL

- a) Tablespoon amounts (1/4 to 1/2 ounce) of bait should be placed at 8 - 12 foot intervals.
- b) Bait should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes.
- c) An uninterrupted supply of bait should be maintained for at least 15 days.

NOTE Bait stations should be checked regularly and refilled before empty. Baits containing PIVALYL CONCENTRATE have excellent keeping qualities and are highly acceptable to rats and mice. If baits are moldy, soured, or infested with insects, they should be replaced.

CAUTION: PIVALYL CONCENTRATE is an anti-coagulant, which if ingested may reduce the clotting ability of the blood. PIVALYL CONCENTRATE and baits prepared with it should carefully be guarded against feeding by humans, poultry, livestock, pets and domestic animals. If, by chance the bait or concentrate is ingested, give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

Do Not Reuse Empty Container. Destroy It By Burying With Waste Or Burning. Stay Away From Smoke Or Fumes.

NOTE FOR PHYSICIANS: Intravenous injection of Vitamin K₁ is indicated together with administration of blood transfusions.

WARNING: BAIT SHOULD BE PLACED IN LOCATIONS NOT ACCESSIBLE TO CHILDREN, PETS, AND DOMESTIC ANIMALS OR IN TAMPER-PROOF BAIT BOXES. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. **WEIGHT;**

MANUFACTURED BY **YORK CHEMICAL COMPANY, Inc.** GARDEN CITY PARK, N. Y.

12/1/77
1748-17