



CERTOX
POWER IN
EVERY DROT

Code Number R-20 Plus

PROLIN* READY-TO-USE RAT & MOUSE BAIT

ACTIVE INGREDIENTS

Sulfaquinoxaline (N-(2-quinoxalinyll (sulfanilamide) (sulfaquinoxaline)	0.025%
Warfarin 3-(3-allyloxybenzyl)- 4-hydroxycoumarin	0.025%

INERT INGREDIENTS

99.950%

TOTAL 100.000%

USDA REG. NO. 1748-78

Net Weight 1 lb. or 453.60 grams

CAUTION:

KEEP OUT OF REACH OF CHILDREN!

Read carefully caution statement on back panel

GENERAL DIRECTIONS

- Place baits in locations where rats and mice feed, water, or travel.
- Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable.
- Contaminated or spoiled baits should be replaced promptly.
- Baiting should continue until all signs of feeding have stopped.

DIRECTIONS FOR RAT CONTROL

- Four ounces of bait per station should be the MINIMUM amount used.
- Bait should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes.
- An uninterrupted supply of bait should be maintained for at least 15 days.

DIRECTIONS FOR MOUSE CONTROL

- Tablespoon amounts (1/4 to 1/2 ounce) of bait should be placed at 8-12 foot intervals.
- Bait should be placed in locations not accessible to children, pets and domestic animals, or in tamper-proof bait boxes.
- An uninterrupted supply of bait should be maintained for at least 15 days.

CAUTION:

Protect children, pets, wildlife and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

CAUTION: In case of poisoning, give 1 tablespoonful of salt in a glass of warm water. Repeat until vomit fluid is clear.

Note to Physician: When a human has been known to have ingested the Meal-baited transfusion combined with intravenous injection and oral doses of Vitamin K are indicated as in the case of hemorrhage by overdoses of Dicumarol.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

CERTOX CHEMICAL COMPANY

Garden City Park, L. I., N. Y.

Printed in U. S. A.

Made in U. S. A.