

certox

BAIT

R
O
Z
O
L

york

Only Rozol Baits are actually impregnated
by the Toxicant

Active Ingredient: 2-[(p-chlorophenyl) phenyl- acetyl] - 1,3-indandione*	0.005%
Inert Ingredients:	99.995%
	100.00%

*Chlorophacinone-Liphadione—Patented in the U.S.A.

Keep away from humans, domestic animals and pets. If swallowed or absorbed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In that case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Call a physician immediately. Baits must be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes.

DO NOT REUSE CONTAINER
DESTROY WHEN EMPTY!
EPA Reg. No. 1748-127
EPA Est. 1748-NY-1

RAT &
MOUSE
BAIT

READY
TO USE

inc.

Place baits in locations where rats and mice feed, water, and/or travel. Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable. Contaminated or spoiled baits should be replaced promptly.

Protect bait against access of children, pets, livestock and wildlife.

FOR NORWAY AND ROOF RATS: Each station should contain at least 4 ounces of bait. Maintain constant supply of bait at stations for at least 10 days.

FOR HOUSE MICE: Tablespoon amounts (¼ to ½ ounce) of bait should be placed 8 to 12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

KEEP AWAY FROM FOODSTUFFS

Sold by: **YORK CHEMICAL CO., INC.**

195 ATLANTIC AVENUE
Garden City Park, New York 11040