Ampicide 9

ACTIVE INGREDIENTS	
Disodium cyanodithioimidocarbonate	[illegible]
Potassium N-methyldithiocarbamate	[illegible]
INERT INGREDIENTS	[illegible]
TOTAL	[illegible]

This product contains [illegible] per gallon

KEEP OUT OF REACH OF CHILDREN

WARNING

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING [illegible] Wear [illegible] Avoid contact [illegible]

STATEMENT OF PRACTICAL TREATMENT [illegible]

ENVIRONMENTAL HAZARDS [illegible]

[illegible]

[illegible] STORAGE AND DISPOSAL

[illegible] product [illegible] Ampicide 9 [illegible] 1000 gallons of water is recommended. Repeat total dosage [illegible] as needed.

[illegible] shock additions of 2.3 to 13.9 fluid ounces of Ampicide 9 per 1000 gallons of water should be employed every 1 to 5 days [illegible] as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the [illegible] shock additions should be made in the sump of [illegible]

STORAGE AND DISPOSAL

PROHIBITIONS [illegible]

PESTICIDE DISPOSAL [illegible]

CONTAINER DISPOSAL [illegible]

GENERAL [illegible]

[illegible]