

DUST OR SPRAY

LUCKY SEVIN Beetle Buster kills a wide range of insects both by contact and through the stomach. Apply with a good duster as insects appear. Treat both upper and lower parts of foliage. Repeat at 7 to 10 day intervals or as needed. In general 1-lb. will treat a garden approx. 25' x 25'. **FOR WET SPRAY** USE mix 1-lb. in 5 gal. of water, or about 10 tbs. per gal. of water. Make a slurry by mixing a small amount of water with Sevin then add water to make desired gallonage. Keep agitated while spraying & spray upper & lower portions of all foliage thoroughly.

CROPS	INSECTS CONTROLLED	AMOUNT TO USE PER ACRE.	APPLICATION
BEANS	Mexican Bean Beetles.	25 lbs. per acre.	Apply when insects appear, then as needed. May be applied up to & including day of harvest. Do not feed treated bean vines to dairy or meat animals.
Cabbage	Leafhopper, Flea Beetle, Bean leaf beetle.	40 lbs. per acre.	Apply when insects first appear, then as needed. May be applied up to and including day of harvest.
TOMATOES EGGPLANT PEPPERS CUCUMBERS SUMMER SQUASH	Flea beetle, Horn worm, Pickleworm, fruit worm, melon worm, corn borer, fall army worm, cucumber beetles, squash bug, leafhoppers.		Apply when insects first appear, then as needed. May be applied up to and including day of harvest.
POTATOES	Cabbage Caterpillar	30 lbs. per acre.	do not apply within 3 days of harvest.
	Colorado potato beetle.	30 to 40 lbs. per acre.	Apply when insects first appear, then as needed. May be applied up to and including day of harvest.
	Leafhoppers, flea beetles, lygus bugs.	25 lbs. per acre.	
CORN	Corn earworm, European corn borer, Japanese Beetle, Fall army worm.	30 to 40 lbs. per acre.	Apply at 2 to 3 day intervals starting when first silk appears, and continue until silk begins to dry. 3 or more applications may be needed, depending on lengths of silking, and severity of insects. Timing & Coverage essential for best control. On sweet corn, may be applied to day of harvest. Fodder & Forage, do not apply within 7 days of Harvest.
Peas	Pea Weevil	30 to 50 lbs.	(Pea Weevil) application should be made at intervals of 4 to 5 days from the time the peas start blossoming until harvest.
	Pea Aphid	40 to 50 lbs. PER ACRE.	Apply when insects first appear. Repeat as needed.
ORNAMENTALS AND FLOWERS such as: Roses, Azalea, Gardenias, etc.	Japanese Beetle, Bagworm, Aphid, Blister Beetles, Boxelder bug, Certain Leaf miners, Elm Leaf Beetle, Flea Beetles, Leafhopper, Leaf rollers, orange tortrix, Periodical cicada, scale insects, Tent caterpillars and exposed thrips, Birch leaf miner.		



BEAN


BEETLE

Vegetable Dust

BETTER
BUSTER

100

FORMULA NO.



USDA REG. NO. 1767-15

STRAWBERRIES

Meadow spittlebug & Strawberry leaf roller.

30 to 40 lbs. per acre.

Apply in post bloom & summer sprays, repeat application as insects appear. May be applied up to one day of harvest.

ACTIVE INGREDIENTS:

Rotenone

Other Cube Resins

Carbaryl- 1-naphthyl N-methylcarbamate

INERT INGREDIENTS:

75%
1.50%
3.00%
94.75%

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe



KEEP OUT OF REACH OF CHILDREN
USE ONLY FOR GARDEN PURPOSES

ORNAMENTALS
AND
FLOWERS such
as: Roses
-zalea
radio, etc.

Japanese Beetle, Bagworm, Aphid, Blister Beetles, Boxelder bug, Certain Leaf miners, Elm Leaf Beetle, Flea Beetles, Leafhopper, Leaf rollers, orange tortrix, Periodical cicada, scale insects, Tent caterpillars and exposed thrips, Birch leaf miner.


until harvest.
Apply when insects first appear. Repeat as needed.



USDA REG. NO. 1767-15

FORMULA NO.

100



NEAT VET

BEAN

BEETLE

For

Fruits

Flowers

Vegetables

Ornamentals

Vegetable Dust

BEE-TLE
BUSTER

KEEP OUT OF REACH OF CHILDREN
SEE INSTRUCTIONS FOR ADULT USE

STRAWBERRIES

Meadow spittlebug & Strawberry leaf roller.

30 to 40 lbs. per acre.

Apply in post bloom & summer sprays, repeat application as insects appear. May be applied up to one day of harvest.

ACTIVE INGREDIENTS:

Rotenone

Other Cube Resins

Carbaryl- 1-naphthyl N-methylcarbamate

INERT INGREDIENTS:

.75%

1.50%

3.00%

94.75%

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

Do not apply to vegetable crops within 1 day of harvest.

On cucumbers, summer squash and peas 1 pound treats a garden approx. 40' x 40'

