

Do not use sulphur on apple foliage within ten days to two weeks of an oil spray.

Sulphur may burn foliage when temperature is high. Do not make sulphur applications at such times.

FOR SPRAYING: Use at the rate of 3 1/2 level tablespoonfuls in 1 gallon of water, or 4 pounds in 100 gallons. TO MIX: stir the required amount of material into the water, and in a power sprayer add it to the water in the tank with the agitator running. When using it in a hand sprayer, shake frequently to improve the agitation.

FOR DUSTING: Use material as is in a good dust gun. Dust lightly and thoroughly from both sides of plants when little air is in motion.

THIS MATERIAL may be used with LEAD ARSENATE if desired.

NON-WARRANTY—Since the use of the product is beyond the control of the Seller, the Buyer assumes all risks of use or handling, whether in accordance with directions or not, as the Seller makes no warranty, expressed or implied, concerning this product.

WILL LOWMY

LUCKY
STRIKE

MADE IN U.S.A.

SULPHUR FOR DUSTING AND SPRAYING

FOR BLACK SPOT, POWDERY MILDEW ON ROSES,
BROWN ROT OF STONE FRUITS, SCAB ON APPLES,
PEARS, PEACHES and certain other plant
diseases. Also in control of Red Spiders on
flowers and ornamentals.



FOR FRUIT TREES

REGISTERED

ACCEPTED

FEB 11 1967

UNDER THE FEDERAL INSECTicide
FUNGICIDE AND RODENTICIDE ACT
FOR CONTROL OF PESTS REGISTERED
TO UNDER NO. 167-28 SUBJECT
TO ANNUAL COMMENTS.

CAUTION KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

Apply Lucky Strike Sulphur as a dust or spray for the control of such diseases as Black Spot of Roses, Powdery Mildew of Roses, and on Zinnias, Sweet Peas and Lilacs, Rust on Snapdragon and Carnations, Brown Rot of Stone Fruits, Scab on Apples, Pears and Peaches, and to aid in keeping Red Spiders in check on evergreens, and where they are present on flowers or other ornamentals. Do not use on evergreen until after new growth has hardened.

APPLY every week or ten days during the growing season on flowers and ornamentals, or more often during rainy weather. Frequent light dust or spray applications are better than heavy applications. Cover all parts of the plant with a thin film of dust or spray. On stone fruits it is a good practice for Brown Rot control to apply this material just as the trees are coming into bloom, and after the petals fall, and at intervals of two weeks during the season. During the last month before harvest dust stone fruits lightly every week or ten days. Delay applications on evergreens until the new growth has hardened. For control of scab on apples and pears make weekly applications from the time new growth appears in the Spring until 30 days after petal fall. Use more frequently during rainy periods.

1 LB. NET WT.