

Manufactured by
THE PARROTT CHEMICAL CO.
Stamford, Connecticut

CAUTION

HARMFUL IF SWALLOWED

If taken internally call a physician **IMMEDIATELY**. Until he arrives follow the directions here given. **ANTIDOTE:** Induce vomiting by tickling the throat with a feather or the finger and give an emetic of strong mustard water, or if available, use Syrup of Ipecac, 1 to 2 tablespoons, then give lime water, milk and honey, white of egg and mucilaginous drinks in copious draughts.

Caution: This produce is not recommended in early rainy seasons when temperature is below 50 degrees F. and not on Stone Fruits such as Peaches or Plums in the growing period.

ACTIVE INGREDIENT:

Copper (in Basic Copper Sulfate)

Expressed as Metallic

INERT INGREDIENTS

12.75%

87.25%

USD, REG. NO. 1767-28

FORMULA NO
1100

**LUCKY
STRIKE**

1 LB NET WT

**BLIGHT
MIXTURE**

12.75% METALLIC COPPER

Used for the same blight & disease conditions on crops as
Bordeaux Mixture.

**FOR USE AS
A BLIGHT AND DISEASE
PREVENTATIVE MATERIAL
ON POTATOES, TOMATOES,
CUCUMBERS AND CERTAIN
OTHER CROPS.**

**CAUTION KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS**

DIRECTIONS

Unless otherwise stated begin applications before diseases appear and repeat at 7 to 10 day intervals. For use on the following crops mix 12 to 16 level tablespoonfuls to each gallon of water, or 1 lb. to 4 or 5 gallons of water. Mix 10 to 15 lbs. to 50 gallons of water per acre. Dust at the rate of 25 to 50 lbs. per acre.

POTATOES; Early & Late Blight, **TOMATOES;** Early & Late Blight, Septoria leaf spot, anthracnose, leaf mold, **CARROTS;** Leaf Spot & Early & Late Blight, **BEANS;** Downy Mildew, **CELERY;** Early & Late Blight, Leaf Spot, **BEETS, PEPPERS, EGG PLANTS;** Leaf Spot, Early and Late Blight.

For use on the following crops mix 6 to 8 level tablespoonfuls to each gallon of water or 1/2 lb. to 4 or 5 gallons of water. Mix 5 to 7 lbs. to 30 gallons of water per acre. Dust at rate of 8 to 15 lbs. per acre.

CUCUMBERS, SQUASH, MELONS, CANTALOUPE; Anthracnose, Downy Mildew, Angular Leaf Spot, Cabbage, Broccoli, **CAULIFLOWER;** Downy Mildew, Leaf Spot. **STRAWBERRIES;** Leaf Spot. **GRAPES;** Black rot, anthracnose, Downy mildew. **ORNAMENTAL PLANTS & SHRUBS;** Black Spot.

When dusting, use a good duster, dust lightly and thoroughly. On grapes & strawberries do not apply within two weeks of Harvest.

This is a general guide for average conditions. As local conditions vary we suggest you consult State Agricultural Experiment station or State Extension Service Authority.