

1 LB. NET WT.

GRAPES AND PLUMS

INSECTS - Leafhopper, Spider Mites, Mealybugs, Mealy Plum Aphid. Apply before insects first appear. Generally begin application in the pre-bloom stage and repeat as needed, usually 10 to 14 day intervals.

Do not treat grapes within 14 days of harvest.

STRAWBERRIES

INSECTS - Red Spider Mite, Spittle Bug, Strawberry Weevil, Leafhopper, Leaf Roller.

DISEASES - Botrytis Fruit Rot.

Apply before insects or disease appears, and repeat as needed. Generally begin application in pre-bloom period and repeat as needed.

Do not treat strawberries within 14 days of harvest

CAUTION:

Wash edible portions of fruit after harvest to remove any remaining residue. Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust or spray mist. Avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and warm water after use. Avoid contamination of food and foodstuffs. Keep out of reach of children and domestic animals.

Malathion upon repeated or prolonged use may cause cholinesterase inhibition. If poisoning occurs, get prompt medical attention. Atropine is antidotal.

CAUTION:

The use of this material at a time or in a manner other than in accordance with the directions may produce plant injury. Do not apply when temperature is above 90° F. Do not apply to fruits during the full bloom period. Do not apply within 7 days of harvest.

ORNAMENTALS, FLOWERS, SHRUBS AND SMALL TREES

INSECTS - Dog Worms, Box Wood Leaf Miner, Oak Kermites, Tent Caterpillar, Azalea Scale, Black Scale Crawlers, Euonymus Scale, Flat Scale, Florida Red Scale, Pear Scale, Leaf Scale, Magnolia Scale, Japanese Scale, Oyster Shell Scale, Scaly Scale, Soft Brown Scale, Scale Insects, Aphids, Birch Leaf Miner, Four-lined Leaf Bug, Japanese Scale, Lace Bugs, Leaf Hoppers, Mealybugs, Tarnished Plant Bug, European Elm Shoot Moth.

DISEASE - Black Spot Disease.

Apply before insects or disease appears, and repeat as needed.

The use of the material being beyond our knowledge and control and involving elements of risk to vegetation, we make no warranty, expressed or implied, as to the effects of such use, whether or not in accordance with directions or claims so to be.

Active Ingredients:

Captan*	2.00%
Malathion**	1.00%
Methoxychlor (Technical)***	8.00%
Inert Ingredients	89.00%
Total	100.00%


*N-trichloromethylthiotetrahydrophthalimide.
**O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.
***Equivalent to 7.4% of 2,2-bis (p-methoxyphenyl) -1,1,1-trichloroethane and 0.96% other isomers and related compounds.

Caution

This product is toxic to fish

Manufactured by THE DUNLOP CHEMICAL COMPANY, STANFORD, CALIF.


CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE BACK PANEL FOR ADDITIONAL CAUTIONS.




TRADE MARK REG. U.S. PAT. OFF.

TRADE MARK REG. U.S. PAT. OFF.

4888



USDA REG. NO. 1760



DIRECTIONS FOR USE

The directions below are general. For exact timing of spray or dust consult your State Experimental Station. Further applications of other insecticides or fungicides may be required for specific pests in your area. This product will not replace the dormant spray on fruit.

APPLES AND PEARS

INSECTS - Aphids, Bud Moth, Forster Scale, Codling Moth, Plum Curculio, Pear-Scale, Pear Leaf Miner, Spider Mites.

DISEASES - Apple Scab, Tree's Spot, Fly Speck, Bitter Rot, Black Rot, Crisp Rot, Botrytis.

In general, application should begin when fruit buds are in delayed dormant stage of development (small leaves beginning to appear) and repeated at 10 to 14 day intervals except during bloom. Do not apply when temperature is above 90° F. to prevent foliage injury.


PEACHES

INSECTS - Spider Mites, Oriental Fruit Moth, Plum Curculio, Japanese Pear Aphid.

DISEASES - Peach Scab, Brown Rot.

Apply before insects or disease first appear, and repeat as needed. Generally begin first application in the pre-pink stage of bud development (when petals show pink). May cause off flavor on peaches. Repeat at 10 to 14 day intervals.

Do not treat peaches within 21 days of harvest.



SPRAYING: Mix at the rate of 6 level tablespoons to each gallon of water or 1-1/2 to 8 gallons of water. Keep well agitated while spraying. Apply as a fine mist spray and cover all parts of the foliage and fruit thoroughly. DUSTING: Apply with a good duster to all parts of fruit and foliage. Apply evenly. Excessively heavy applications may cause injury.

CHERRIES

INSECTS - Black Cherry Aphid, Fruit Tree Leaf Roller.

TOMATOES

INSECTS - Aphids, Spider Mites, Tomato Fruit Worm.
DISEASES - Early Blight, Late Blight, Septoria, Stemphylium, Anthracnose. Apply before insects or disease appears, and repeat as needed.

CUCUMBERS, SQUASH AND MELONS

INSECTS - Aphids, Spider Mites, Striped Cucumber Beetle, 12 Spotted Beetle, Pickleworm, Melonworm, Squash Vine Borer.
DISEASES - Angular Leaf Spot, Anthracnose, Downy Mildew. Apply before insects or disease appears, and repeat as needed.

POTATOES

INSECTS - Aphids, Leafhoppers.
DISEASES - Early Blight, Late Blight. Apply before insects or disease appears, and repeat as needed.

NOTE: Do not apply to Fruits or Vegetables later than 7 days before harvest. Do not apply to cucurbits unless plants are dry.

Fruits, Vegetables, Flowers, Ornamentals
KILLS & CONTROLS MANY
DANGEROUS PESTS

1 LB. NET WT.

GRAPES AND PLUMS

INSECTS — Leafhopper, Spider Mites, Mealybugs, Mealy Plum Aphid.
Apply before insects first appear. Generally begin application in the pre-bloom stage and repeat as needed, usually 10 to 14 day intervals.

Do not treat grapes within 14 days of harvest.

STRAWBERRIES

INSECTS — Red Spider Mite, Spittle Bug, Strawberry Weevil, Leafhopper, Leaf Roller.

DISEASES — Botrytis Fruit Rots.

Apply before insects or disease appears, and repeat as needed. Generally begin application in pre-bloom period and repeat as needed.

Do not treat strawberries within 14 days of harvest

CAUTION:

Wash edible portions of fruit after harvest to remove any remaining residue. Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust or spray mist. Avoid getting in eyes, on skin, or on clothing. Wash thoroughly with soap and warm water after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals.

Malathion upon repeated or prolonged use may cause cholinesterase inhibition. If poisoning occurs, get prompt medical attention. Atropine is antidotal.

CAUTION:

The use of this material at a time or in a manner other than in accordance with the directions may produce plant injury. Do not apply when temperature is above 90° F. Do not apply to fruits during the full bloom period. Do not apply within 7 days of harvest.

ORNAMENTALS, FLOWERS, SHRUBS AND SHADE TREES

INSECTS — Bag Worms, Box Wood Leaf Miner, Oak Kermes, Tent Caterpillar, Azalea Scale, Black Scale Crawlers, Euonymus Scale, Fletcher Scale, Florida Red Scale, Juniper Scale, Leaf Scale, Magnolia Scale, Monterey Pine Scale, Oyster Shell Scale, Scurfy Scale, Soft Brown Scale, Spider Mites, Aphids, Birch Leaf Miner, Four-lined Leaf Bug, Japanese Beetle (adult), Lace Bugs, Leaf Hoppers, Mealybugs, Tarnished Plant Bug, Thrips, Whitefly, European Pine Shoot Moth.

DISEASE — Black Spot Disease.

Apply before insects or disease appears, and repeat as needed.

The use of the material being beyond our knowledge and control and involving elements of risk to vegetation, we make no warranty, expressed or implied, as to the effects of such use, whether or not in accordance with directions or claims so to be.

Active Ingredients:

Captan*-----	8.00%
Malathion**-----	4.00%
Methoxychlor (Technical)***-----	8.00%
Inert Ingredients-----	80.00%
Total	100.00%

*N-trichloromethylthiotetrahydrophthalimide.

**O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

***Equivalent to 7.4% of 2,2-bis (p-methoxyphenyl) -1,1,1-trichloroethane and 0.96% other isomers and related compounds.

Caution

This product is toxic to fish

Manufactured by THE PARROTT CHEMICAL COMPANY, STAMFORD, CONN.


CAUTION KEEP OFF SEE BACK PANEL FOR

STAMFORD

TRADE MARK

4888


USDA REG.



The use of this material at a time or in a manner other than in accordance with the directions may produce plant injury. Do not apply when temperature is above 90° F. Do not apply to fruits during the full bloom period. Do not apply within 7 days of harvest.

Caution
This product is toxic to fish

Manufactured by THE PARROTT CHEMICAL COMPANY, STAMFORD, CONN.




FRUIT TREES

MAIL CAP
TRADE MARK REG. U.S. PAT. OFF.
FOR

4888



USDA REG. NO. 1767-000



DIRECTIONS FOR USE

The directions below are general. For exact timing of spray or dust consult your State Experimental Station. Further applications of other insecticides or fungicides may be required for specific pests in your area. This product will not replace the dormant spray on fruit.

APPLES AND PEARS

INSECTS — Aphids, Bud Moth, Forbes Scale, Codling moth, Plum Curculio, Red Banded Leaf Roller, Spider Mites.

DISEASES — Apple Scab, Crook's Spot, Fly Spock, Bitter Rot, Black Rot, Black Pox, Botryosphaeria.

In general, application should begin when fruit buds are in delayed dormant stage of development (small leaves beginning to appear) and repeated at 10 to 14 day intervals except during bloom. Do not apply when temperature is above 90° F. to prevent foliage injury.

PEACHES

INSECTS — Spider Mites, Oriental Fruit Moth, Plum Curculio, Japanese Beetle, Aphids.

TO MIX AND APPLY

SPRAYING: Mix at the rate of 6 level tablespoons to each gallon of water or 1-lb. to 8 gallons of water. Keep well agitated while spraying. Apply as a fine mist spray and cover all parts of the foliage and fruit thoroughly. **DUSTING:** Apply with a good duster to all parts of fruit and foliage. Apply evenly. Excessively heavy applications may cause injury.

CHERRIES


INSECTS — Black Cherry Aphid, Fruit Tree Leaf Roller.

TOMATOES

INSECTS — Aphids, Spider Mites, Tomato Fruit Worm. **DISEASES** — Early Blight, Late Blight, Septoria, Stemphylium, Anthracnose. Apply before insects or disease appears, and repeat as needed.

CUCUMBERS, SQUASH AND MELONS

INSECTS — Aphids, Spider Mites, Striped Cucumber Beetle, 12 Spotted Beetle, Pickleworm, Melonworm, Squash Vine Borer. **DISEASES** — Angular Leaf Spot, Anthracnose, Downy Mildew. Apply before insects or disease appears, and repeat as needed.



DIRECTIONS FOR USE

The directions below are general. For exact timing of spray or dust consult your State Experimental Station. Further applications of other insecticides or fungicides may be required for specific pests in your area. This product will not replace the dormant spray on fruit.

APPLES AND PEARS

INSECTS — Aphids, Bud Moth, Forbes Scale, Codling moth, Plum Curculio, Red Banded Leaf Roller, Spider Mites.

DISEASES — Apple Scab, Brook's Spot, Fly Speck, Bitter Rot, Black Rot, Black Pox, Botryosphaeria.

In general, application should begin when fruit buds are in delayed dormant stage of development (small leaves beginning to appear) and repeated at 10 to 14 day intervals except during bloom. Do not apply when temperature is above 90° F. to prevent foliage injury.

PEACHES

INSECTS — Spider Mites, Oriental Fruit Moth, Plum Curculio, Japanese Beetle, Aphids.

DISEASES — Peach Scab, Brown Rot.

Apply before insects or disease first appear, and repeat as needed. Generally make first application in the pre-pink stage of bud development (when bud first shows pink). May cause off flavor on peaches. Repeat at 10 to 14 day intervals.

Do not treat peaches within 21 days of harvest.

APPLES	TOMATOES	ROSES
PEACHES	SQUASH	
PEARS - PLUMS	CUCUMBERS	
STRAWBERRIES	GRAPES	

TO MIX AND APPLY

SPRAYING: Mix at the rate of 6 level tablespoons to each gallon of water or 1-lb. to 8 gallons of water. Keep well agitated while spraying. Apply as a fine mist spray and cover all parts of the foliage and fruit thoroughly.
DUSTING: Apply with a good duster to all parts of fruit and foliage. Apply evenly. Excessively heavy applications may cause injury.

CHERRIES

INSECTS — Black Cherry Aphid, Fruit Tree Leaf Roller.

TOMATOES

INSECTS — Aphids, Spider Mites, Tomato Fruit Worm.

DISEASES — Early Blight, Late Blight, Septoria, Stomphylium, Anthracnose. Apply before insects or disease appears, and repeat as needed.

CUCUMBERS, SQUASH AND MELONS

INSECTS — Aphids, Spider Mites, Striped Cucumber Beetle, 12 Spotted Beetle, Pickleworm, Melonworm, Squash Vine Borer.

DISEASES — Angular Leaf Spot, Anthracnose, Downy Mildew. Apply before insects or disease appears, and repeat as needed.

POTATOES

INSECTS — Aphids, Leafhoppers.

DISEASES — Early Blight, Late Blight.

Apply before insects or disease appears, and repeat as needed.

NOTE: Do not apply to Fruits or Vegetables later than 7 days before harvest. Do not apply to cucurbits unless plants are dry.

Fruits, Vegetables, Flowers, Ornamentals
KILLS & CONTROLS MANY
INSECTS & DISEASES