

LAWNS and AROUND DWELLINGS — Against Clover Mites (Brown Mite, Clover Spider Mite or brown Almanca Mite) Mix 2 tps. per 1 gal. of water. Spray lower foundations of house as well as ground, lawns and plants in 10 ft. wide area along side of house. Repeat as needed. Do not use inside dwellings.

FLIES, around Animal Quarters, Outbuildings, Picnic Areas, and Dwellings — Mix 5 tbs. per 1 gal. of water. Adding 3 to 5 tbs. of cane or dark sugar or molasses may increase effectiveness. Spray around house foundations, under porches, or other fly breeding places. 1 gal. diluted spray treats area 10 ft. x 10 ft. Repeat as needed. Remove animals from buildings before treatment. **CAUTION** — Do not use in milk rooms.

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. AVOID INHALATION. DO NOT GET IN EYES, SKIN OR CLOTHING. WASH THOROUGHLY WITH SOAP AND WATER AFTER USE. CHANGE CLOTHING IMMEDIATELY AFTER USE AND WASH BEFORE RE-USE. DESTROY CONTAINERS WHEN EMPTY. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. KEEP FROM CHILDREN AND DOMESTIC ANIMALS. MALATHION UPON REPEATED OR PROLONGED USE MAY CAUSE CHOLINESTERASE INHIBITION. IF POISONING OCCURS, GET PROMPT MEDICAL ATTENTION. Atropine IS ANTIDOTAL. USE carefully.

ACTIVE INGREDIENTS

MALATHION*

Xylol

INERT INGREDIENTS

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

E.P.A. REG.

50%

40% NO. 1767- 44

10%

This product is toxic to fish. Keep out of lakes, streams and ponds

FOR

FRUIT TREES
FLOWERS

50%

MALATHION

Emulsifiable

WITH TRITON X-190

Lucky
Strike

FORMULA
NO. 99

KILLS: APHIDS & MITES,
BEETLES
RED SPIDER FLIES JAP BEETLE
MEALY BUGS & SCALE
CAUTION KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

ONE PINT (16 FL. OZ.)

BEFORE USING READ DIRECTIONS AND NOTE CAUTIONS

Apply Spray thoroughly on both upper and lower leaf surfaces and other infested plant parts. Repeat treatment as needed.

FRUITS — On APPLES, PEACHES, PEARS, CHERRIES, GRAPES, STRAWBERRIES, and PLUMS, against Aphids, Mix 1½ to 2 tps. to 1 gal. water. Against Red Spider Mites, Mealybugs, Woolly Aphis, White Fly, Thrips, Tarnished Plant Bug, Four-Lined Leaf Bug, Pear Psyllid, Japanese Beetle Adults, Codling Moth, Fruit Tree Leaf Roller, Plum Curculio and Red Banded Leaf Roller, Mix 1 tbs. per gal. water. Against SCALE CRAWLERS mix 1½ to 2 tbs. per gal. water. Do not apply to Apples, Cherries, Plums, Strawberries within 10 days of harvest. Do not apply to Grapes, Peaches, or Pears within 14 days of harvest. Against Plum Curculio and Codling Moth make applications with full coverage timed in accordance with your State Agricultural Experiment station recommendations.

VEGETABLES — COLE CROPS, BEANS, BEETS, BROCCOLI, CABBAGE, LETTUCE, PEAS, PEPPERS, POTATOES, TOMATOES, against Aphids, Mix 1½ to 2 tps. to 1 gal. water. Against Red Spider Mites, Mealybugs, White Fly, Thrips, Tarnished Plant Bug, Four-Lined Leaf Bug, Bean Leafhopper, Potato Leafhopper, Japanese Beetle Adults, Cabbage Looper, Mexican Bean Beetle, Imported Cabbage Worm, Mix 1 tbs. per 1 gal. water. Do not apply to Peas within 14 days of harvest. Do not apply to other vegetables within 10 days of harvest.

FLOWERS, ORNAMENTAL TREES and SHRUBS, EVERGREENS — ROSES, CAMELIAS, AZALEAS, CHRYSANTHEMUMS, GLADIOLUS, ASTERS, LILACS, GREENHOUSE PLANTS, PHLOX, ARBOR VITAE, BOXWOOD, PINES, SPRUCES, VINES, YEW, against Aphids, Lacebug, 1½ to 2 tps. to 1 gal. of water. Against Red Spider, Mites, Mealybugs, Woolly Aphis, White Fly, Thrips, Tarnished Plant Bug, Four-Lined Bug, Rose Leafhopper,

1467-44

LAWNS and AROUND DWELLINGS — Against Clover Mites (Brown Mite, Clover Spider Mite or Brown Almond Mite) Mix 2 tsps. per 1 gal. of water. Spray lower foundations of house as well as ground, lawns and plants in 10 ft. wide area along side of house. Repeat as needed. Do not use inside dwellings.

FLIES, around Animal Quarters, Outbuildings, Picnic Areas, and Dwellings — Mix 5 tbs. to 1 gal. of water. Adding 3 to 5 tbs. of cane or dark sugar or molasses may increase effectiveness. Spray around house foundations, under porches, or other fly breeding places. 1 gal. diluted spray treats area 10 ft. x 10 ft. Repeat as needed. Remove animals from buildings before treatment. **CAUTION** — Do not use in milk rooms.

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. AVOID INHALATION. DO NOT GET IN EYES, SKIN OR CLOTHING. WASH THOROUGHLY WITH SOAP AND WATER AFTER USE. CHANGE CLOTHING IMMEDIATELY AFTER USE AND WASH BEFORE RE-USE. DESTROY CONTAINERS WHEN EMPTY. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. KEEP FROM CHILDREN AND DOMESTIC ANIMALS. MALATHION UPON REPEATED OR PROLONGED USE MAY CAUSE CHOLINESTERASE INHIBITION. IF POISONING OCCURS, GET PROMPT MEDICAL ATTENTION. Atropine IS ANTIDOTAL. USE carefully.

ACTIVE INGREDIENTS

MALATHION*

Xylol

INERT INGREDIENTS

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

E.P.A. REG.
50%
40% **NO. 1767- 44**
10%

This product is toxic to fish. Keep out of lakes, streams and ponds.

BEFORE USING READ DIRECTIONS AND NOTE CAUTIONS

Apply Spray thoroughly on both upper and lower leaf surfaces and other infested plant parts. Repeat treatment as needed.

FRUITS — On ~~APPLES, PEACHES, PEARS, CHERRIES, GRAPES, STRAWBERRIES, and PLUMS,~~ against Aphids, Mix 1½ to 2 tsps. to 1 gal. water. Against Red Spider Mites, Mealybugs, Woolly Aphis, White Fly, Thrips, Tarnished Plant Bug, Four-Lined Leaf Bug, Pear Psyllid, Japanese Beetle Adults, Codling Moth, Fruit Tree Leaf Roller, Plum Curculio and Red Banded Leaf Roller, Mix 1 tbs. per gal. water. Against SCALE CRAWLERS mix 1½ to 2 tbs. per gal. water. Do not apply to Apples, Cherries, Plums, Strawberries within 10 days of harvest. Do not apply to Grapes, Peaches, or Pears within 14 days of harvest. Against Plum Curculio and Codling Moth make applications with full coverage timed in accordance with your State Agricultural Experiment station recommendations.

VEGETABLES — ~~COLE CROPS, BEANS, BEETS, BROCCOLI, CABBAGE, LETTUCE, PEAS, PEPPERS, POTATOES, TOMATOES,~~ against Aphids, Mix 1½ to 2 tsps. to 1 gal. water. Against Red Spider Mites, Mealybugs, White Fly, Thrips, Tarnished Plant Bug, Four-Lined Leaf Bug, Bean Leafhopper, Potato Leafhopper, Japanese Beetle Adults, Cabbage Looper, Mexican Bean Beetle, Imported Cabbage Worm, Mix 1 tbs. per 1 gal. water. Do not apply to Peas within 14 days of harvest. Do not apply to other vegetables within 10 days of harvest.

FLOWERS, ORNAMENTAL TREES and SHRUBS, EVERGREENS — ~~ROSES, CAMELIAS, AZALEAS, CHRYSANTHEMUMS, GLADIOLUS, ASTERS, LILACS, GREENHOUSE PLANTS, PHLOX, ARBOR VITAE, BOXWOOD, PINES, SPRUCES, VINES, YEW,~~ against Aphids, Lacebug, 1½ to 2 tsps. to 1 gal. of water. Against Red Spider, Mites, Mealybugs, Woolly Aphis, White Fly, Thrips, Tarnished Plant Bug, Four-Lined Bug, Rose Leafhopper, Japanese Beetles Adults, European Pine Shoot Moth, Mix 1 tbs. per 1 gal. of water. Against Bagworm, Birch Leafminer, Boxwood Leafminer, Tent Caterpillar, Mix 1 to 1½ tbs. per gal. of water. Against Scale Crawlers, Mix 1½ to 2 tbs. per gal. of water. Do not use on certain ferns including Boston, Maidenhair and Pteris, and some species of Crassula.

Mfd. by **THE PARROTT CHEMICAL COMPANY, Stamford, Conn.**