

GRAPES: To control Grape Leaf Miner, Grape Leafhopper, Grape Berry Moth, Japanese Beetle, apply when insects first appear, then as needed. Do not apply within 2 weeks of harvest. Do not use stickers after backset of fruit.

DOWNY MILDEW: Apply before disease first appears and repeat every 7 to 10 days. Do not apply within 2 weeks of harvest.

Apply as directed to control downy mildew by spraying or using other effective means. Apply the correct amount as indicated on label.

CAUTION: Avoid inhalation. Wash face and hands after using. Avoid prolonged or repeated skin contact. Get away from product if contact is made.

CAUTION: Keep out of lakes, streams and ponds, do not contaminate water by cleaning of equipment or disposal of wastes. Dispose of containers and wastes by burying them in crop lands away from water supplies. Do not use as fertilizer.

ACTIVE INGREDIENTS:	
(Carbaryl) 1-naphthyl N-methylcarbamate	5.00%
Copper (Basic Copper sulphate)	7.00%
Expressed as Metallic	
INERT INGREDIENTS	88.00%
TOTAL	100.00%

Manufactured by THE PARROTT CHEMICAL CO., STAMFORD, Conn.

● **CAUTION** KEEP OUT OF REACH OF CHILDREN

5% SEVIN

7% COPPER METALLIC

BLIGHT

CONTROLS: EARLY & LATE

POTATO-MAKER

KILLS CERTAIN INSECTS ON

POTATOES TOMATOES

GRAPES

DIRECTIONS

1 LB NET WT.

USDA REG. NO. 1767-75

LUCKY STRIKE

FORMULA NO. 400

Apply with a good dusting or insect spray. Treat both upper and lower parts of foliage. Repeat at 7 to 10 day intervals until needed. In general 1-lb. will treat a garden approx. 25' x 25'. For wet spray use mix 1-lb. in 5 gal. of water, or about 10 to 15 per gal. of water. Make a slurry by mixing a small amount of water then add water to make desired gallonage. Keep at hand while spraying and spray upper and lower portions of all foliage thoroughly.

POTATOES: To control Colorado Potato Beetle, Leafhoppers, Flea Beetles, Lygus bugs, apply when insects first appear then as needed. May be applied up to and including day of harvest.

EARLY AND LATE BLIGHT: Apply before disease appears and repeat every 7 to 10 days up to and including day of harvest. Do not feed cooked potatoes to dairy animals.

TOMATOES: To control late blight and worms, apply when insects first appear, then as needed. May be applied up to and including day of harvest.

EARLY and LATE BLIGHT, SEPTORIA LEAF SPOT, ANTHRACNOSE, LEAF MOLD: Apply before disease appears and repeat every 7 to 10 days up to and including day of harvest.