

DIRECTIONS FOR MIXING

Mix at the rate of 3 tablespoonsfull per gallon of water. MALTOX is both an insecticide and fungicide for use in all conventional type sprayers and garden hose spray guns.

DIRECTIONS FOR USE

Spray thoroughly upper and lower sides of foliage when plants begin to grow and repeat at 7 to 14 day intervals, or as needed.

FLOWERS & ORNAMENTAL S: Spray Roses, Dahlias, Asters, Camelias, Chrysanthemums, Gladioli, for control of Japanese Beetles, Aphids, Thrips, Leaf Hoppers, Flea Beetles, Mealy Bug, Spittle Bugs, Plant Bugs, Lace Bugs, Cankerworms, White Flies and Black Spot on Roses.

SHADE TREES AND SHRUBS: Spray Shade Trees, Evergreens, Azaleas, Spirea, Lilacs, for control of Red Spider mites, Bird Damage and Film Leaf Miners, Soft Brown Scale, Black Scale Crawlers, Oyster Shell Scale, Cankerworms, Tent caterpillars, Leaf Roller Caterpillars, Gypsy Moth Worm (Larvae), Gypsy Moth Adult.

VEGETABLES: TOMATOES, For control of Aphids and Late Blight. Do not apply within 7 days of harvest. **BEANS,** for control of Mexican Bean Beetle, Leaf Hoppers and Anthraco nose. Do not apply within 7 days of harvest.

FRUIT TREES: Apples, Spray apples against Red Banded Leaf Roller, Codling Moth, Spider Mites, Tent Caterpillars, Apple Scab, Bitter Rot apply at 7 to 10 day intervals. Do not use within 7 days of harvest.

CHERRIES, Spray against Red Banded Leaf Roller, Aphids, Brown Rot, Blossom Blight apply at 7 to 10 day intervals. Do not use within 7 days of harvest. **PEACHES,** Spray against Spider Mites, Oriental Fruit Moth, Aphids, Brown Rot, Blossom Blight, apply at 7 to 10 day intervals. Do not use on peaches within 21 days of harvest.

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid inhalation of spray mist. Do not get in eyes, skin or clothing. Wash thoroughly with soap and water after use. Change clothing immediately after use and wash before re-use. Wear protective clothing. May cause skin irritation. Avoid contamination of feed and foodstuffs. Keep from children and domestic animals. Malathion upon use may cause cholinesterase inhibition. If poisoning occurs get prompt medical attention; Atropine is antidotal.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective Information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on label. Destroy containers when empty.

SEE BACK PANEL FOR ADDITIONAL INFORMATION

GARDEN FERTILIZER

FOR USE IN ALL TYPES

CAUTION: KEEP OUT OF REACH OF CHILDREN

Malathion

SPRAY

MALTOX

INSECTICIDE



ONE PINT (16 FL OZ)


not apply within 7 days of harvest. **BEANS**, for control of Mexican Bean Beetle, Leaf Hoppers and Anthracnose. Do not apply within 7 days of harvest.

FRUIT TREES: Apples, Spray apples against Red Banded Leaf Roller, Codling Moth, Spider Mites, Tent Caterpillars, Apple Scab, Bitter Rot apply at 7 to 10 day intervals. Do not use within 7 days of harvest. **CHERRIES**, Spray against Red Banded Leaf Roller, Aphids, Brown Rot, Blossom Blight apply at 7 to 10 day intervals. Do not use within 7 days of harvest. **PEACHES**, Spray against Spider Mites, Oriental Fruit Moth, Aphids, Brown Rot, Blossom Blight, apply at 7 to 10 day intervals. Do not use on peaches within 21 days of harvest.

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid inhalation of spray mist. Do not get in eyes, skin or clothing. Wash thoroughly with soap and water after use. Change clothing immediately after use and wash before re-use. Wear protective clothing. May cause skin irritation. Avoid contamination of feed and foodstuffs. Keep from children and domestic animals. Malathion upon use may cause cholinesterase inhibition. If poisoning occurs get prompt medical attention; Atropine is antidotal.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on label. Destroy containers when empty.



*O, O-DIMETHYL DITHIOPHOSPHATE OF DIETHYL MERCAPTOSUCCINATE.
** EQUIVALENT TO 17.6% OF 2,2-BIS (P-METHOXYPHENYL)-1, 1, 1 -
TRICHLOROETHANE AND 2.4% OTHER ISOMERS AND RELATED COMPOUNDS.
*** 2,3 - DICHLORO-1, 4 - NAPHTHOQUINONE.