

BEANS

Lacebug, Tomato fruit worm, ~~PUMPKIN SQUASH, CUCUMBER, MELON~~. Flea beetle, Melonworms, Cucumber beetle, Squash bug. Avoid excess applications. Some leaf injury may occur if treatments are made when foliage is wet or in presence of prolonged high humidity. Do not use on watermelons in Florida.

CAUTIONS: HARMFUL IF INHALED OR SWALLOWED: Avoid breathing of spray. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands and face before eating. Avoid contamination of food and keep out of reach of children. Do not re-use container. Destroy when empty. Do not store below 32°F. Do not freeze.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, Cholinesterase Inhibitor. Atropine is Antidotal.

E.P.A. REG. NO. 1767-84

TRADE MARK OF UNION CARBIDE CORP.

PACKED BY THE PARROTT CHEMICAL COMPANY
STAMFORD, CONN. 06904

KILLS: GYPSY MOTH, CHINCH BUGS, JAPANESE BEETLES, TENT CATERpillARS, BAGWORM AND CERTAIN OTHER ORNAMENTAL, LAWN AND GARDEN INSECTS.

ACTIVE INGREDIENTS

CARBARYL 1-NAPHTHYLENEMETHYL CARBAMATE 42.0%

INERT INGREDIENTS 57.4%

CAUTION KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

FORMULA
NO. 42

ONE PINT MAKES 50 GALLONS FINISHED SPRAY



ONE PINT (16 FL. OZ.)

LIQUID FLOWABLE
SEVIN

KEEP COVER ON OPENED CONTAINER TO PREVENT DRYING

DIRECTIONS: Flowable Sevin is an extremely fine dispersion of Sevin in water. SHAKE WELL BEFORE USING. Pour recommended amount of material into nearly full spray tank or pre-mix material in another container half full of water and pour mixture into spray tank. Add balance of water. Agitate while mixing and using. Apply as a coarse wet spray to thoroughly cover all plant surfaces.

PRECAUTIONS: Avoid spray drifting to adjoining food or forage crops or crops in bloom. For protection of honey bees, avoid unnecessary use during periods when honey bees are visiting crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application. Consult state or county agricultural extension authorities for timing and uses consistent with label cautions and directions. Sevin is compatible with common insecticides, fungicides and miticides, but not with lime or other highly alkaline materials.

LAWN SPRAY: For Ants, Chinch Bugs, Earwigs, Fall armyworm, Fleas, Mosquitoes, Sod webworms (lawn moths) and Millipedes. For best results treat after cutting and when the ground is moist from rain or irrigation using 1/4 pint in 15 to 20 gallons of water to each 500 square feet of established lawn. Apply with hose end or pressure sprayer with ample water to insure turf penetration. Repeat in 2 to 3 weeks if necessary.

ORNAMENTAL PLANT SPRAY: For flowering shrubs and Shade Trees including roses, azaleas, mums, lilac, privet, dogwood, magnolia, oak and maple. (Do not use on Boston Ivy). Apply spray mixture containing 2 teaspoonfuls per gallon of water (4 fl. oz. to 12 gals. or 1 pint to 50 gals.) when insects or damage appear. Repeat at weekly intervals or as necessary for such insects as apple aphid, bagworm, leaf miners, elm leaf beetle, gypsy moth, Japanese beetle, lacebug, rose slug, tent caterpillars, leaf rollers, scale insects and periodical Cicada.

VEGETABLE GARDEN SPRAY: Mix 1 fl. oz. in 3 gallons of water (1 quart in sufficient water for 1 acre). Spray when insects or damage appear and repeat at 7 to 10 day intervals when necessary. No post treatment time limitations for the following host plants when material is applied in accordance with directions.