

Directions for Use

Annual Weeds:

Apply one gallon of VEG-OUT (diluted with up to 10 gallons of water) per 150 to 200 square feet when plants are young. Thoroughly wet with a sprayer or sprinkling can, all parts of the plant, particularly the growing tips.

Perennial Weeds:

For the best results treat plants during their dormant stage or when nearing maturity. Good results are obtained with Spring application, but Summer applications are not recommended. Apply by spraying 1 to 2 gallons of VEG-OUT per 100 square feet. For deep rooted weeds and grass, such as Bermuda, Johnson, Bindweed and Quack grass the heavier rate is recommended. Retreatment may be necessary to secure adequate control of certain hard-to-kill weeds.

Note: This material is non-selective and effects all types of vegetation. Extreme care should be exercised in its use to prevent killing lawns, shrubs, trees and other valuable plants. This includes the treatment of areas which may be overlaid by tree roots.

CAUTION: HARMFUL IF SWALLOWED. Avoid contamination of feed and foodstuffs. Keep away from children and domestic animals. (Do not contaminate domestic or irrigation water.)

Do not reuse empty container. Wash thoroughly with water and detergent. Discard in safe place.

NOTICE

KEEP FROM FREEZING. If this product is stored in cold or freezing weather it may separate or thicken. This will not harm the performance. Before using, warm to room temperature and stir thoroughly.

PRINTED IN U.S.A.

NATIONAL CHEMSEARCH CORPORATION

Research Laboratory and Plant

1000 Rte. 100, Suite 100

Edison, N.J.

State Line, Monmouth County

New Jersey

Western Plant

1000 Rte. 100, Suite 100

Edison, N.J.

CAUTION:

KEEP OUT OF REACH OF CHILDREN

(See Side Panel for additional cautions.)

NATIONAL

CHEMSEARCH

VEG-OUT

NON-SELECTIVE


Bermuda grass, Poison Ivy, Poison Oak, Bindweed, Canada wildflower, and other grasses and weeds. Toxic to all types of vegetation. Used for vegetation control on non-cropped areas such as around buildings, along roadsides, power lines, right-of-ways.

Active Ingredients

Sodium Chlorate 18.00%
Sodium Metaborate 6.00%

Inert Ingredients:

76.00%



NATIONAL

CENTEX™

ACCEPTED
June 13, 1975
 1769-82
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO.

NON-SELECTIVE

ACTIVE INGREDIENTS:

Sodium Chlorate 18.00%
 Sodium Metaborate 9.00%
 INERT INGREDIENTS 73.00%

CAUTION:

KEEP OUT OF REACH OF CHILDREN
 (See right panel for additional precautions.)

EPA Reg. No. 1769—82—AA

NET CONTENTS: _____

NATIONAL CHEMSEARCH 2727 CHEMSEARCH BLVD. IRVING, TEXAS 75062
 Eastern Plant: 1000 N. Main St., Memphis, Tenn. Western Plant: 4500 N. 10th St., Phoenix, Ariz.
 NATIONAL CHEMSEARCH, INC. Copyright © 1975 National Chemsearch, Inc.
 DALLAS ST. LOUIS HACKENSACK BEVERLY HILLS INDIANAPOLIS

CONTROLS:

Johnson Grass, Quack Grass, Bermuda Grass, Poison Ivy, Poison Oak, Bindweed, Canada Thistle, Sand Burr and many other grasses and weeds. Toxic to all type of vegetation. This material is recommended for vegetation control on non-cropped areas such as fence-rows, ditch banks, around buildings, along roadsides, power lines, rights-of-way and other similar waste areas.

DIRECTIONS FOR USE

Annual Weeds:

Apply 1 gallon of CENTEX (diluted with up to 10 gallons of water) per 150 to 200 square feet when plants are young. Thoroughly wet with a sprayer or sprinkling can all parts of the plant, particularly the growing tips.

Perennial Weeds:

For the best results, treat plants during their dormant stage or when nearing maturity. Good results are obtained with Spring application, but Summer applications are not recommended. Apply by spraying 1 to 2 gallons of CENTEX per 100 square feet. For deep rooted weeds and grass such as Bermuda, Johnson, Bindweed and Quack grass, the heavier rate is recommended. Re-treatment may be necessary to secure adequate control of certain hard to kill weeds.

FOR INDUSTRIAL OR
 INSTITUTIONAL USE ONLY

Note: This material is non-selective and affects all types of vegetation. Extreme care should be exercised in its use to prevent killing lawns, shrubs, trees and other valuable plants. This includes the treatment of areas which may be underlaid by tree roots.

Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

CAUTION: Avoid contamination of feed and foodstuffs. Keep away from children and domestic animals. Do not contaminate water used for humans, domestic animal consumption or for irrigation. Harmful if swallowed.

NOTICE

KEEP FROM FREEZING. If this product is stored in cold or freezing weather it may separate or thicken. This will not harm the performance. Before using, warm to room temperature and stir thoroughly.

ENVIRONMENTAL CAUTIONS: Do not contaminate domestic or irrigation water by cleaning of equipment or disposal of wastes. **APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.**

4-75

EPA Est. No. 4140

CENTEX INDUSTRIAL

CERTIFIED LABORATORIES

APTREX

Non-Selective Weed Killer - Soil Sterilant *

* Semi-permanent soil sterilant toxic to all forms of plant life.

Will kill most weeds and grasses including Bermuda, Johnson Grass and Dollar Grass.
Will also suppress nut grass.

Use around industrial sites, lumber yards, boundary fences, warehouses, bridges,
sign posts, fire hydrants, utility poles, power substations, outdoor theaters,
tennis courts, school playgrounds and parks.

ACTIVE INGREDIENTS:

Sodium Chlorate	12.7%
*Sodium Metaborate	7.0%
Bromacil (5-bromo-3-sec-butyl- 6-methyluracil)	.4%

INERT INGREDIENTS:

79.9%

*Equivalent to 1.16% boron expressed as elemental boron

CAUTION: (18 pt. tp.) KEEP OUT OF REACH OF CHILDREN (12 pt. tp.)
(See right panel for additional caution)

DIRECTIONS: In areas of light rainfall (under 20 inches annual) mix 7 quarts of
APTREX with sufficient water to cover 1000 square feet. Apply in such a manner
to get uniform coverage.

In areas of over 20 inches of rainfall annually use 3 1/2 gallons of APTREX in
sufficient water to thoroughly cover 1000 square feet.

Time of Application - For best results apply in the spring when weeds and grasses
are small and growing vigorously. In areas of low annual rainfall, earlier
applications may be desirable.

NOTE: This material is non-selective and effects all types of vegetation. Extreme
care should be exercised in its use to prevent killing lawns, shrubs, trees and
other valuable plants. This includes the treatment of areas which may be underlain
by tree roots. Due to the varied conditions of weather, soil, type of vegetation,
location and time of applications, the manufacturer does not guarantee complete kill
of vegetation or accept responsibility for its use or mis-use.

CAUTION: Avoid contamination of feed and foodstuffs. Keep away from all
domestic animals. HAZARDOUS IF SWALLOWED.

U.S.D.A. P. C. NO. 1

Approved by the Department of
Agriculture, U.S. Department of
Agriculture, U.S. Department of
Agriculture, U.S. Department of

