

This product is highly toxic to bees exposed to direct treatment  
or residues on crops. Protective informa-  
tion may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish and wildlife.  
Keep out of lakes, streams & ponds.  
Apply this product only as specified on label.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS.

Do not apply when weather conditions favor drift of  
spray from areas treated

# DIRECTIONS

BORER	TREES OR SHRUBS	DOSE	TIME TO APPLY
PINE BARK BEETLES	Pines	1 pint/5 gals. water	Spray trunks thoroughly - wet trunks to run-off.
BARK BEETLES, FLAT HEAD BORERS, ROUND HEAD BORERS	Felled Logs, Rough Timber	1 pint/5 gals. water	As soon as trees are felled, spray all surface areas thoroughly. Wet to run-off.
PINE ROOT COLLAR WEEVIL, WHITE PINE WEEVIL	Pine, Spruce	3 pints/100 gals. water 1 Tablespoonful/gal. water	Spray lower trunks and around base of trees, soaking bark and wetting soil surfaces. White Pine Weevil - Apply late April when buds begin to swell. Spray upper parts of trees thoroughly.
TAXUS WEEVIL, STRAWBERRY ROOT WEEVIL	Taxus, Hemlock	1 Tablespoonful / 1 gal. water	Apply in May or early June to the trunks and around base of trees soaking the bark and wetting the soil surface. Repeat in July for Strawberry Root Weevil adult control.
FRUIT TREE BARK BEETLES, FLAT-HEADED BORERS, ROUND-HEADED BORERS	Apples, Cherries, Peaches	1 Tablespoonful / 1 gal. water 3 pints/100 gals. water	Apply to bark of tree trunk and large limbs in spring. Apply to trunk, crotch to ground. Soak bark to run-off. Do not apply to fruit or foliage.
BARK BEETLES, WOOD BORING BEETLES	Maple, Aspen, Poplar, Willows, Elm, Locust	3 pints/50 gals. water 2 Tablespoonfuls/1 gal. water	Apply in spring when buds begin to break. Repeat in 10-14 days. Elms - During summer, repeat in fall - soak trunks to run-off.
PEACH TREE BORER	Peaches	3 pints/100 gals. water 1 Tablespoonful/gal. water	Spray lower trunk and around base of tree, soaking bark and wetting trunk from crotch to ground. Do not apply to fruit or foliage.
DOGWOOD BORER, RHODODENDRON BORER, LILAC BORER	Dogwood, Rhododendron, Lilac	1 Tablespoonful/ 1 gal. water	Make 2 to 3 applications 7 to 10 days apart in May and early June.
LEAF MINERS	English and Scotch Elms, Birch, Locust	1 Teaspoonful to 1 gallon water	Apply when small miners appear in early and mid May Apply when small miners appear in early July

CAUTION: Harmful if swallowed. Avoid inhalation and skin contact. May be absorbed through skin. In case of contact, wash immediately with soap and water. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Keep out of reach of children. Do not use or store near heat or open flame. When container is empty, immediately wash thoroughly and destroy. Never re-use.

MANUFACTURED BY THE PARROTT CHEMICAL CO., STAMFORD, CONN.

ACCEPTED  
10-2-72  
1767-69