

FOR ECONOMIC POISON REGISTERED UNDER NO. 1769-113



C-400

CRAB GRASS KILLER



Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and bury in a safe place.

KEEP OUT OF REACH OF CHILDREN

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

CERTIFIED C-400 CRAB GRASS KILLER is to be used for destroying Crab Grass in established turf except Centipede, Carpet and St. Augustine grasses.

ACTIVE INGREDIENTS:

Dodecylammonium methanearsonate 8.0% *
Octylammonium methanearsonate 8.0% *

INERT INGREDIENTS: 84.0%

* Total arsenic, calculated as elemental arsenic, all in water soluble form - 4.1%

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near food and feed products. Store away from heat. Keep children and animals off of treated areas until washed into soil.

KEEP OUT OF REACH OF CHILDREN

YOU CAN RELY ON



An exclusive product of CERTIFIED LABORATORIES, INC.
Ft. Worth, Tex. • Chicago, Ill. • Kendall Park, N.J.
Mountain View, Calif.
P. O. BOX 2493 FT. WORTH, TEXAS 76101

C-400 in the seeding or early branched stage, however, it will control all stages. For best results, good soil moisture is essential. When applied during period of drought, a thorough watering of the area is necessary prior to application. Keep area mowed at from 1" to 1½" during treatment. New lawns or areas should not be treated until after 3 mowings. Wanted grasses may be temporarily affected, especially during hot, dry weather. Even in sensitive Bent and Fescue grasses may be temporarily discolored. Thorough watering will usually revive them. The more persistent Bluegrasses, Zoysia, Bermuda and Dichondra are not usually affected. Do not use on Centipede, Carpet or St. Augustine grass.

DIRECTIONS: For each 1000 square feet of area, apply 5 to 8 ounces of C-400. Mix in 5 gallons of water when temperature is above 85° F. When temperature is below 85° F, use 6 1/2 to 8 ounces to 5 gallons of water. Thoroughly wet all crab grass. For best control, two applications five to seven days apart is generally required. Repeat treatment as necessary.

For Dallis Grass: Apply at the rate of 2 to 4 gallons of C-400, mixed with 100 gallons of water per acre. Repeat two or more times every 5 to 7 days. Observe the same precautions as for Crab Grass control.

USDA Reg. No. 1769-113

10-67

PRINTED IN U.S.A.

Nov 20, 1967
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 1769-113

KEEP OUT OF REACH OF CHILDREN

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

CAUTION: Harmful if absorbed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near food and feed products. Store away from heat. Keep children and pets out of treated areas until dry.

CGK-79 Crab Grass Killer is selected for the selective control of crab grass with little or no injury to good lawn grasses except St. Augustine, Carpet and Centipede grass. CGK-79 Crab Grass Killer should be applied when the crab grass is young, preferably from the seedling to the early branched stage. Early treatment is preferred although CGK-79 Crab Grass Killer controls both types. Good soil moisture will give best results. During dry spells, water thoroughly the area to be treated before applications are made. Keep the area mowed at about 1" to 1 1/2" during treatment. On new lawns or areas,



NATIONAL CHEMSEARCH CORPORATION
 Research Laboratory and Plant
 2730 Carl Road, Irving, Texas
 Eastern Plant: Stouts Lane,
 Monmouth Junction, New Jersey
 Western Plant: 932 Kifer Road,
 Sunnyvale, California

CAUTION:
 KEEP OUT OF REACH OF CHILDREN

**CGK-79
 CRAB GRASS KILLER**

ACTIVE INGREDIENTS:
 Dichloromethyl p-dimethylaminoacetate 20.0%
 Octylamine 20.0%
 INERT INGREDIENTS 60.0%
 Total weight calculated as 100% based on 100% active ingredient.
 All in water-soluble form.

