

National Chemsearch ACTI-CIL can be used to kill most weeds for an extended period of time on non-cropland areas such as loading ramps, fence rows, railroad sidings, storage yards, parking lots, around buildings and industrial plant sites. Apply ACTI-CIL, undiluted, 1 gallon per 1000 square feet.

Some of the weeds controlled by ACTI-CIL are foxtail, ryegrass, wildoats, crabgrass, cheatgrass, broomsedge, ragweeds, lambsquarter and turkey mullein. Johnson grass, Salt grass, Bouncing Bet, Brackern fern, Horsetail, Smooth Broome, Canada Bluegrass, purpletop, quackgrass, aster dandelion, dog fennel, goldenrod, plantain and wildcarrot.

NOTE: Do not let the liquid come in contact with or spray to drift on desirable plants. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. Do not store near fertilizers, seeds, insecticides, or fungicides. Do not contaminate water used for irrigation or domestic purposes.

CAUTION: Harmful if swallowed or absorbed through skin. Causes irritation to skin. Do not breathe spray mist or vapor. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. In case of contact wash skin with soap and water. For eyes, immediately flush with large amounts of water for at least 15 minutes and get medical attention. Remove and wash contaminated clothing before reuse.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Apply this product only as specified on this label. Do not contaminate water by cleaning off equipment or disposal of wastes.

PRINTED IN U.S.A.

ACTI-CIL

WEED KILLER

KILLS MOST VEGETATION

ACTIVE INGREDIENTS


Heavy Aromatic Naphtha	50.00%
Petroleum Base Oil	46.51
Bromacil (5-bromo-3-sec-butyl-6-methyl-uracil)	0.60%
Pentachlorophenol	0.80%
Other chlorophenols	0.09%

INERT INGREDIENTS 2.00%

CAUTION:

KEEP OUT OF REACH OF CHILDREN

(See Side Panel for additional cautions.)



National Chemsearch ACTI-CIL can be used to kill most weeds for an extended period of time on non-cropland areas such as loading ramps, fence rows, railroad sidings, storage yards, parking lots, around buildings and industrial plant sites. Apply ACTI-CIL, undiluted, 1 gallon per 1000 square feet.

Some of the weeds controlled by ACTI-CIL are foxtail, ryegrass, wildoats, crabgrass, cheatgrass, broomsedge, ragweeds, lambsquarter and turkey mullein. Johnson grass, Salt grass, Bouncing Bet, Brackern fern, Horsetail, Smooth Broome, Canada Bluegrass, purpletop, quackgrass, aster dandelion, dog fennel, goldenrod, plantain and wildcarrot.

NOTE: Do not let the liquid come in contact with or spray to drift on desirable plants. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. Do not store near fertilizers, seeds, insecticides, or fungicides. Do not contaminate water used for irrigation or domestic purposes.

CAUTION: Harmful if swallowed or absorbed through skin. Causes irritation to skin. Do not breathe spray mist or vapor. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. In case of contact wash skin with soap and water. For eyes, immediately flush with large amounts of water for at least 15 minutes and get medical attention. Remove and wash contaminated clothing before reuse.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Apply this product only as specified on this label. Do not contaminate water by cleaning off equipment or disposal of wastes.

USDA Reg. No. 148-111-110-09