

[1769-229]

**ACTIVE INGREDIENTS:**

*+(5-Benzyl-3-furyl) methyl 2,2-dimethyl-2-(2-methylpropenyl) cyclopropanecarboxylate . . . . .	0.250%
Related compounds . . . . .	0.034%
Aromatic petroleum hydrocarbons . . . . .	0.331%
Petroleum distillate . . . . .	19.375%
<b>INERT INGREDIENTS:</b> . . . . .	<b>80.010%</b>

\*Cis/trans isomers ratio: max. 30% (+) cis and min. 70% (+) trans.

+U.S. Patent No. 3,465,007

**CAUTION:** KEEP OUT OF REACH OF CHILDREN. Avoid contamination of feed and foodstuffs. Cover or remove fishbowls. Use as a space spray in food processing plants only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. In hospitals, patients should be removed from room prior to treatment. Room should be ventilated for 2 hours after spraying. Do not return patients to room until after ventilation.

**WARNING:** Contents under pressure. Do not puncture. Do not expose container to temperatures above 100°. Never throw container into fire or incinerator.



**DIRECTIONS FOR USE:**

Remove protective cap. Aim spray opening away from person. Push button to spray.

**FOR HOUSEFLIES, MOSQUITOES, AND GNATS:** Close all doors and windows. Point nozzle upward, direct spray mist to all parts of room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

**WASPS:** Application should be made in late evening when insects are at nest. Spray liberally into hiding and nesting places, contacting as many insects as possible. Repeat as necessary.

**FOR CRAWLING INSECTS:** Roach, earwigs, grass mites, sowbugs, silver and centipedes. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects where possible.

**ANTS:** Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

**CAUTION:** Do not use in food preparation areas.

**ACTIVE INGREDIENTS:**

- \*+(5-Benzyl-3-furyl) methyl 2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate . . . . .0.250%
- Related compounds . . . . .0.034%
- Aromatic petroleum hydrocarbons . . .0.331%
- Petroleum distillate . . . . .19.375%

**INERT INGREDIENTS:** . . . . .80.010%

\*Cis/trans isomers ratio: max. 30% (+) cis and min. 70% (+) trans.

+U.S. Patent No. 3,465,007

**CAUTION:** KEEP OUT OF REACH OF CHILDREN. Avoid contamination of feed and foodstuffs. Cover or remove fishbowls. Use as a space spray in food processing plants only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. In hospitals, patients should be removed from room prior to treatment. Room should be ventilated for 2 hours after spraying. Do not return patients to room until after ventilation.

**WARNING:** Contents under pressure. Do not puncture. Do not expose container to temperatures above 120°F. Never throw container into fire or incinerator.



Pressurized Spray Insecticide

For ~~Bakeries, Food Plants,~~  
~~Restaurants, Schools, Hotels,~~  
~~Motels, Office Buildings,~~  
~~Hospitals and Food~~  
~~Handling Plants.~~

KEEP OUT OF REACH OF CHILDREN

Net Weight 11.0 oz.

**DIRECTIONS FOR USE:**

Remove protective cap. Aim spray opening away from person. Push button to spray.

**FOR HOUSEFLIES, MOSQUITOES, AND GNATS:** Close all doors and windows. Point nozzle upward, direct spray mist to all parts of room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

**WASPS:** Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary.

**FOR CRAWLING INSECTS:** ~~Roaches,~~ earwigs, grain mites, sowbugs, spiders and centipedes: Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insect where possible.

**ANTS:** Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

**NATIONAL CHEMSEARCH CORPORATION**  
 DALLAS • ST. LOUIS • NEW YORK • LOS ANGELES • TORONTO  
 Research Laboratory and Plant  
 2730 Carl Road Irving, Texas 75061