

LEFT PANEL

NATIONAL CHEMSEARCH

PRETROL

Selective Pre-Emergent Weed Killer

A selective pre-emergence herbicide for the control of annual grasses and certain broadleaf weeds in ornamental turf.

ACTIVE INGREDIENTS:

*Dimethyl tetrachloroterephthalate.....6.0%

INERT INGREDIENTS:.....94.0%

TOTAL.....100.0%

* Use as a pre-emergent herbicide patented under U.S. Patent No.

2,923,634

This product contains the toxic inert ingredient formaldehyde.

EPA Reg. No. 1769-246

CAUTION:

KEEP OUT OF REACH OF CHILDREN

See side panel for additional precautions.

NET CONTENTS:

National Chemsearch
Div. of NCH Corporation
2727 Chemsearch Blvd.
Irving, Texas 75062

ACCEPTED
DEC 7 1967

1769-246

CENTER PANEL

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Do not breathe spray mist. Avoid contact with skin, eyes, or clothing. In case of contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS:

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Apply this pesticide only as specified on the label.

2064

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician immediately.

STORAGE AND DISPOSAL:

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: KEEP CONTAINER CLOSED.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

PLASTIC CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of the smoke.

METAL CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

NOT FOR RESALE

RIGHT PANEL

BEST AVAILABLE COPY

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do Not Apply This Product Through Any Type Of Irrigation System.

For the control of some grasses and broadleaf weeds in established turfgrass such as smooth and hairy crabgrass, green foxtail, yellow foxtail, lovegrass, witchgrass, carpetweed, common chickweed, common lambsquarters, purslane, Florida pusley, apply one gallon of National Chemsearch PRETRON in five to

ten gallons of water to cover 2,000 square feet.

One application, applied before weeds germinate in early spring, usually provides season long control. Crabgrass seed germination usually coincides with the time when Forsythia blossoms start to fall. In areas which commonly experience stands of late-germinating crabgrass and other annual weedy grasses, a second application at half the regular rate can be made safely two months after the first application. When spring seeding is necessary, there should be a delay of approximately 60 days after applying PRETROL.

Early spring application of PRETROL may be applied to newly seeded turf after it exhibits greening of newly sprouted turfgrass. Early spring applications of PRETROL allow fall sowing of permanent grass.

Poa Annuua - To control annual bluegrass and other late germinating grasses and weeds, a late summer or early fall application of 1 1/2 gallons of National Chemsearch PRETROL in 10 gallons of water per 2,000 square feet should supplement the early spring application. Earlier applications are required in Southern States. Consult agricultural experiment station or extensive service weed specialist for date Poa Annuua seed germinates in your area and make application at least two weeks prior to date.

To control redroot, pigweed, sandbur, dodder, barnyardgrass, goosegrass, broun top panicum, ground cherry, johnson grass (from seed), nodding spurge, spotted spurge, and prostate spurge. An early spring application of 1 1/2 gallons of PRETROL in 10 gallons of water per 2000 sq. ft. is required.

NOTE: Do not use on Cohansey or Toronto bentgrasses. National Chemsearch PRETROL does not show herbicide injury or discoloration to established common turf grasses. No more than the recommended rate should be used. Do not apply to Dichondra. PRETROL is not recommended for use on bentgrasses mowed at putting green height.

PRETROL is not recommended for use on turf grasses where the following ornamentals are present as possible injury can occur:

Bugleweed
Button pink
Carnation
Geum
Germander
Mesembrythemum
Pansy
Phlox

STIR WELL BEFORE USING AND KEEP AGITATED DURING USE.

BEST AVAILABLE COPY

454

TO REORDER:

In Texas, call 1-800-427-7950

Outside Texas, call 1-800-527-9921

(DATE)

EPA Est. No. 4140

CA-1 IN-1 NJ-1 PR-1 TX-1