

NATIONAL

ORNATHAL

LAWN AND ORNAMENTAL FUNGICIDE

ACTIVE INGREDIENT:

*Chlorothalonil, (tetrachloroisophthalonitrile) 4.5%

INERT INGREDIENTS: 95.5%

*U. S. Patent No. 3,290,353 and 3,331,735

CAUTION:

KEEP OUT OF REACH OF CHILDREN

(See right panel for additional cautions.)

EPA Reg. No. 1769-251-AA

NET CONTENTS:

NATIONAL CHEMSEARCH 2727 CHEMSEARCH BLVD., IRVING, TEXAS 75062
Eastern Plant: Stoughton, Mass. Western Plant: 912 Kifer Rd., Sunnyvale, Calif.
NATIONAL CHEMSEARCH div. of USACHEM Inc. Copyright 1974 National Chemsearch Corp.
DALLAS • ST. LOUIS • HACKENSACK • BEVERLY HILLS PRINTED IN U.S.A.

GENERAL INFORMATION: National Chemsearch ORNATHAL is highly effective for the control of a number of important turf and lawn diseases as well as a number of diseases of ornamentals. Premix ORNATHAL with water in a bucket before introduction into the sprayer tank for final dilution. The tank should be mixed thoroughly and agitation maintained before and during spraying to prevent settling out or clogging of the line or nozzles. This will insure more uniform application.

DIRECTIONS FOR USE:

TURF: Apply ORNATHAL in 5 to 15 gallons of water per 1000 square feet to give uniform coverage according to the following chart:

DISEASE	Rates per 1000 Square Feet	
	Interval of Application	Preventive Curative
Helminthosporium (Leaf Spot & Melting Out)	7 - 10 days	1 qt. 2 qts.
Dollar Spot	7 - 14 days	1 qt. 2 qts.
Brown Patch	7 - 10 days	1 qt. 2 qts.
Gray Leaf Spot	7 - 10 days	1 qt. 2 qts.
Red Thread	7 - 10 days	1 qt. 2 qts.
Stem Rust of Bluegrass	7 - 14 days	2 qts. 2 qts.
Copper Spot	7 - 10 days	2 qts. 2 qts.

Do not mow or water the grass after treatment. Wait until the grass is thoroughly dry, preferably 24 hours after treatment. Do not graze treated areas. Do not feed clippings to live stock. Start spray applications in Spring, before diseases appear and repeat at 7 to 14 days intervals throughout the season. Make applications on a 7 day schedule during periods of wet weather or severe disease conditions. Under severe disease conditions, use the curative rates and apply on a 7 day spray schedule.

FOR INDUSTRIAL OR INSTITUTIONAL USE ONLY

ORNAMENTALS — Apply ORNATHAL at the first sign of disease and repeat at 7- 14-day intervals throughout the season. Make applications on a 7-day schedule during periods of wet weather or severe disease conditions. To control Black Spot, Blossom Blight, use 1 gallon of ORNATHAL to 37 gallons of water. To control Ascochyta Ray Blight and Gray Mold on chrysanthemum, Botrytis Blight on geraniums and iris and Leaf Spot on iris and Powdery Mildew on zinnia, use 1½ gallons of ORNATHAL to 37 gallons of water.

CAUTION: Do not take internally. Avoid contact with eyes. Wash thoroughly with soap and water after handling.

NOTE: The active ingredient may produce temporary allergic side effects characterized by redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. Affected persons respond to treatment with antihistamines or steroid creams and/or systemic steroids. There is no evidence of after effects or permanent injury.

Do not reuse empty container. Return to a drum reconditioner or perforate and crush and bury in a safe place away from water supplies.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

ACCEPTED

FEB 20 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 1769-251

EPA Est. No. 4140

CA 1 IN 1 NJ 1 PR 1 TX 1