

FRONT PANEL

NATIONAL CHEMSEARCH

BANZO 84

CONTROL BROADLEAF WEEDS ON TURFS & LAWNS

ACTIVE INGREDIENTS:

Diethanolamine salt of 2,4-D Dichlorophenoxyacetic Acid*	8.0%
Diethanolamine salt of dicamba (3,6-dichloro-o-anisic Acid)**	4.0%
Diethanolamine salt of Related Acids	0.69%

INERT INGREDIENTS:

87.31%

*Equivalent to 5.42% 2,4-Dichlorophenoxyacetic Acid

**Equivalent to 2.71% 3,6-dichloro-o-anisic acid

CAUTION: KEEP OUT OF REACH OF CHILDREN (See right panel for additional cautions)

EPA REG. NO. _____

NATIONAL CHEMSEARCH, Div.
of USACHEM, Inc.
2727 Chemsearch Blvd.
Irving, Texas 75062

LEFT PANEL

National Chemsearch BANZO 84 was formulated for the control of broadleaf weeds on turfs and lawns, golf fairways, tees, and roughs. May be used on Bermuda, Merion, Kentucky bluegrass, Fescues and Rye grasses.

Weeds Controlled:

Buckhorn	Pigweed
Burdock	Plaintain
Carpet weed	Poison Ivy
Chickory	Poison Oak
Clover	Purslane
Common Chickweed	Ragweed
Curled Dock	Red Clover
Dandelions	Red Sorrel
Dog Fennel	Sheep Sorrel
English Daisy	Stitchwort
Ground Ivy	Speedwell
Hawkweed	Spurge
Henbit	Spurry
Rhewel	White Clover
knotweed	Wild Garlic
Lawn Burweed	Wild Onion
Lambsquarter	Yarrow
Mallow	
Mouse-ear Chickweed	
Morning Glory	
Pepperweed	

Reg.
1769-275

12

LEFT PANEL (continued)

DIRECTIONS FOR USE:

Mix 6 ounces with 2 to 3 gallons of water and apply uniformly to 1000 square feet. Application can be made by a fine spray or watering can. Repeat application after 3 to 4 weeks if hard to kill weeds are present.

Either before or after application, it is wise to properly fertilize the lawn so as to encourage the lawn to heal as rapidly as possible, filling in the bare areas where the weeds have died away. Once the weeds have been removed, the bare areas should be reseeded. Bare areas may be reseeded 3 to 4 weeks after application.

RIGHT PANEL

WHEN TO APPLY: Apply in spring or fall, when weeds are actively growing. Knot-weed is controlled more easily in spring or early summer when weeds are young. Fall germinating weeds may be controlled more effectively when treated in the fall or early winter.

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes and clothing. Avoid inhaling spray mist. In case of skin contact, flush with an abundance of water. For eyes, flush with an abundance of water and get medical attention. Do not take internally. Do not store with foodstuffs. Keep Out of Reach of Children.

APPLICATION CAUTIONS: Avoid drift of spray mist to susceptible plants as this product may injure grapes, tomatoes, beans, peas, other vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. (Coarse sprays are less likely to drift.) Do not apply within the dripline of trees and shrubs. Do not apply when turf is suffering from drought conditions or during very hot weather or damage to grass may result. Do not mow for several days before or after application. Do not apply during periods of heavy rainfall. Do not use on newly seeded areas before at least two mowings. Do not use if you want clover in your turf. In some areas bent grass, carpet, buffalo and St. Augustine grasses may be injured; therefore, do not use on turf where these grasses predominate. Do not transport or store with seeds, fertilizers, insecticides or fungicides. Do not use in or around greenhouses. Store at temperatures above 32°F. If allowed to freeze, thaw and remix before using. Because of danger of contamination, do not use same spray equipment, except for applying other 2,4-D combinations.

ENVIRONMENTAL CAUTIONS: Do not contaminate water intended for irrigation or domestic purposes. Do not pour solution used to clean sprayer near desirable trees or shrubs or where it may contaminate water. When container is empty, immediately wash thoroughly then destroy by crushing or perforating and burying away from water supplies. Do not reuse empty container. Do not use this product for purposes other than those recommended on the label.

Local conditions may affect the use of this chemical. Consult Agricultural Experiment Station of Extension Service Weed specialists for specific recommendations as to dosage and time of application and for control of other weeds.

EPA EST. NO. 1110 CA-1 IN-1 NJ-1 PR-1 TX-1

Reg No.
1769-278