

FRONT PANEL

NATIONAL CHEMSEARCH

BANCO 88

CONTROL BROADLEAF WEEDS ON TURFS & LAWNS

ACTIVE INGREDIENTS:

Diethanolamine salt of 2,4-Dichlorophenoxyacetic Acid\* 8.00%  
Diethylamine salt of dicamba (5,6-dichloro-o-anisic acid)\*\*3.28%  
Diethylamine salt of related Acids 0.53%

INERT INGREDIENTS:

88.19%

\*Equivalent to 5.42% 2,4-Dichlorophenoxyacetic Acid  
\*\*Equivalent to 2.71% 5,6-dichloro-o-anisic acid

CAUTION: KEEP OUT OF REACH OF CHILDREN (See right panel for additional cautions)

EPA REG. NO. \_\_\_\_\_

NATIONAL CHEMSEARCH, Inc.  
of USACHEM, Inc.  
2727 Chemsearch Blvd  
Irving, Texas 75062

LEFT PANEL

National Chemsearch BANCO 88 was formulated for the control of broadleaf weeds on turfs and lawns, golf fairways, tees, and roughs. May be used on Bermuda, Merion, Kentucky bluegrass, fescues and rye grasses.

Weeds Controlled:

Buckhorn	Henbit	Porson Oak
Burdock	Knives	Purslane
Carpet Weed	Knotweed	Ragweed
Chickory	Lawn Burweed	Red Clover
Clover	Lambsquarter	Red Sorrel
Common Chickweed	Mallow	Sheep Sorrel
Crabgrass	Moose-ear Chickweed	Stitchwort
Dandelion	Morning Glory	Speedwell
Dog Fenner	Pepperweed	Spurge
English Daisy	Pigweed	Spurry
Ground Ivy	Plantain	White Clover
Hardweed	Porson Ivy	Wild Garlic
		Wild Onion
		Yarrow

DIRECTIONS FOR USE:

Mix 6 ounce with 2-4 gallon of water and apply uniformly to 1000 square feet. Application can be made by using a sprayer or watering can. Repeat application after 5 to 6 weeks if hard to kill weeds are present.

Either before or after application, it is wise to properly fertilize the lawn so as to encourage the lawn to heal as rapidly as possible, filling in the bare areas.

BEST DOCUMENT AVAILABLE Reg 1767-276

LEFT PANEL (continued)

where the weeds have died away. Once the weeds have been removed, the bare areas should be reseeded. Bare areas may be reseeded 3 to 4 weeks after application.

RIGHT PANEL

**WHEN TO APPLY:** Apply in spring or fall, when weeds are actively growing. Knotweed is controlled more easily in spring or early summer when weeds are young. Fall germinating weeds may be controlled more effectively when treated in the fall or early winter.

**CAUTION:** Harmful if swallowed. Avoid contact with skin, eyes and clothing. Avoid inhaling spray mist. In case of skin contact, flush with an abundance of water. For eyes, flush with an abundance of water and get medical attention. Do not take internally. Do not store with foodstuffs. Keep Out of Reach of Children.

**APPLICATION CAUTIONS:** Avoid drift of spray mist to susceptible plants as this product may injure grapes, tomatoes, beans, peas, other vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. (Coarse sprays are less likely to drift.) Do not apply within the dripline of trees and shrubs. Do not apply when turf is suffering from drought conditions or during very hot weather or damage to grass may result. Do not mow for several days before or after application. Do not apply during periods of heavy rainfall. Do not use on newly seeded areas before at least two mowings. Do not use if you want clover in your turf. In some areas bent grass, carpet, buffalo and St. Augustine grasses may be injured; therefore, do not use on turf where these grasses predominate. Do not transport or store with seeds, fertilizers, insecticides or fungicides. Do not use in or around greenhouses. Store at temperatures above 32°F. If allowed to freeze, thaw and remix before using. Because of danger of contamination, do not use same spray equipment, except for applying other 2,4-D combinations.

**ENVIRONMENTAL CAUTIONS:** Do not contaminate water intended for irrigation or domestic purposes. Do not pour solution used to clean sprayer near desirable trees or shrubs or where it may contaminate water. When container is empty, immediately wash thoroughly then destroy by crushing or perforating and burying away from water supplies. Do not reuse empty container. Do not use this product for purposes other than those recommended on the label.

Local conditions may affect the use of this chemical. Consult Agricultural Experiment Station of Extension Service weed specialists for specific recommendations as to dosage and time of application and for control of other weeds.

EPA EST. NO. 1110 CA-1 IN-1 NJ-1 PR-1 TX-1

*Reg (767-275)*

10