

4

CRUISE AREAS - Facilities where prepared foods are served, such as dining rooms, but excluding areas where food is prepared or held. Mix 1 part DICHLORON II with 11 parts water to control light infestations and 1 part DICHLORON II to 1 part water to quickly reduce heavy infestations. Apply as a spot treatment to selective surfaces such as baseboards, under elements of construction and into cracks and crevices. Avoid treating surfaces likely to be contacted by food. Do not apply when facility is in operation or foods are exposed.

WARNING: - Pests indoors. For Spot Application Only. Do not use in poultry houses or other animal buildings. Do not allow spray to contact food, feedstuffs and water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

OUT-OF-DOOR PESTS: Houseflies - To reduce the annoyance of houseflies, dilute DICHLORON II at 1 gallon to 11 gallons of water and spray outside surfaces of windows, door frames, around garbage cans, trash bins, etc. where houseflies are a problem. Repeat treatment when effectiveness diminishes.

LAWN PESTS. To control armyworms, cutworms, brown dog ticks, chiggers, chinch bugs, crickets, cutworms, earwigs, fleas, grasshoppers and sod webworms, (lawn moths), dilute 1 gallon of DICHLORON II with sufficient water to cover 10,000 sq. ft. Apply as a coarse, low pressure spray using suitable application equipment. Thoroughly water immediately after treatment to wash the insecticide into the turf except for sod webworms. For sod webworms delay watering or mowing the treated area for 12 to 14 hours after treatment. For best results, the lawn should be moist at the time of treatment, and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear; re-treat when necessary. Control (larvae, pupae, hyperodes) weevils: Use DICHLORON diluted in sufficient water for 10,000 sq. ft. Spray suspected problem areas in the spring and again in mid-May. Control Japanese beetle larvae and European spruce sawfly as well as hyperodes when adults are present and actively feeding near surface usually in late May or August. Irrigate treated areas with water, irrigate the treated area with 1 to 2 inch of water to wash the insecticide into the underlying soil.

SPRING AND SUMMER PESTS: Control grasshoppers, ticks, chiggers, crickets, cutworms, earwigs, fleas, grasshoppers, and sod webworms, (lawn moths) and other lawn pests. Dilute 1 gallon of DICHLORON II with sufficient water to cover 10,000 sq. ft. Apply as a coarse, low pressure spray using suitable application equipment. Thoroughly water immediately after treatment to wash the insecticide into the turf except for sod webworms. For sod webworms delay watering or mowing the treated area for 12 to 14 hours after treatment. For best results, the lawn should be moist at the time of treatment, and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear; re-treat when necessary. Control (larvae, pupae, hyperodes) weevils: Use DICHLORON diluted in sufficient water for 10,000 sq. ft. Spray suspected problem areas in the spring and again in mid-May. Control Japanese beetle larvae and European spruce sawfly as well as hyperodes when adults are present and actively feeding near surface usually in late May or August. Irrigate treated areas with water, irrigate the treated area with 1 to 2 inch of water to wash the insecticide into the underlying soil.

SPRING AND SUMMER PESTS: Control grasshoppers, ticks, chiggers, crickets, cutworms, earwigs, fleas, grasshoppers, and sod webworms, (lawn moths) and other lawn pests. Dilute 1 gallon of DICHLORON II with sufficient water to cover 10,000 sq. ft. Apply as a coarse, low pressure spray using suitable application equipment. Thoroughly water immediately after treatment to wash the insecticide into the turf except for sod webworms. For sod webworms delay watering or mowing the treated area for 12 to 14 hours after treatment. For best results, the lawn should be moist at the time of treatment, and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear; re-treat when necessary. Control (larvae, pupae, hyperodes) weevils: Use DICHLORON diluted in sufficient water for 10,000 sq. ft. Spray suspected problem areas in the spring and again in mid-May. Control Japanese beetle larvae and European spruce sawfly as well as hyperodes when adults are present and actively feeding near surface usually in late May or August. Irrigate treated areas with water, irrigate the treated area with 1 to 2 inch of water to wash the insecticide into the underlying soil.

SPRING AND SUMMER PESTS: Control grasshoppers, ticks, chiggers, crickets, cutworms, earwigs, fleas, grasshoppers, and sod webworms, (lawn moths) and other lawn pests. Dilute 1 gallon of DICHLORON II with sufficient water to cover 10,000 sq. ft. Apply as a coarse, low pressure spray using suitable application equipment. Thoroughly water immediately after treatment to wash the insecticide into the turf except for sod webworms. For sod webworms delay watering or mowing the treated area for 12 to 14 hours after treatment. For best results, the lawn should be moist at the time of treatment, and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear; re-treat when necessary. Control (larvae, pupae, hyperodes) weevils: Use DICHLORON diluted in sufficient water for 10,000 sq. ft. Spray suspected problem areas in the spring and again in mid-May. Control Japanese beetle larvae and European spruce sawfly as well as hyperodes when adults are present and actively feeding near surface usually in late May or August. Irrigate treated areas with water, irrigate the treated area with 1 to 2 inch of water to wash the insecticide into the underlying soil.

5

LIGHT PANEL CONT'D

ATTENTION: (Area Control of Ticks and Chiggers): Do not allow public use of treated areas during application or until spray has dried.

MOSQUITOES: To control mosquito larvae, use 1 gallon of DICHLORON II in sufficient water to cover 10 to 20 acres of none to medium vegetation and 5 to 10 acres of medium to heavy vegetation. Apply the spray to provide complete and uniform coverage of infested plants.

To control adult mosquitoes, use 1 gallon of DICHLORON II in sufficient water to cover 10 acres of none to medium vegetation and 5 acres of heavy vegetation.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

TO REORDER:
In Texas, call 1-800-442-7950
Outside Texas, call 1-800-527-9901

EPA Est. No. 4140

CA-1	IN-1	NO-1	PR-1	TX-1
------	------	------	------	------

BEST DOCUMENT AVAILABLE



DURSBAN® BAIT

A Ready To Use Insecticidal Bait For Application To The Soil Surface Of Flower Beds, Vegetable Garden Beds, Lawns, Golf Courses and Other Recreational Turf For Control Of Certain Insects as Listed which may Feed Upon This Insecticidal Bait.

ANTS, SOLE BUGS, FIELD CRICKETS, MOLE CRICKETS, TERMITES,

EARWIGS AND DARKLING BEETLES

one quart treats 100 square feet

ACTIVE INGREDIENT:

Chlorpyrifos [0,0,-Diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate]*..... 0.25%

INERT INGREDIENTS: 99.75%

Total 100.00%

*Dursban Reg. U.S. Dow Chemical Company
U.S. Patent No. 3,244,000

EPA Reg. No. 331-1000

EPA Reg. No. 331-1000-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel For Additional Precautionary Statements

Directions For Practical Treatment

If Swallowed - Call a physician or Poison Control Center immediately. Have victim drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If On Skin - Remove contaminated clothing and wash affected areas with soap and water.

If In Eyes - Flush eyes with water for at least 15 minutes. Get medical attention if irritation persists.

ACCEPTED
with COMMENTS
by EPA Letter

May 1, 1984

100-1000-1
100-1000-1
100-1000-1

It is a white, powdery substance which is effective against a wide variety of insects. It is particularly effective against the following insects:

AS A BAIT AGAINST THE CRICKETS IN THE GARDEN

NOTE

Do not use this product where root crops such as carrots, beets, potatoes, radishes, etc. are grown. The product is poisonous to plants and their parts.

Apply this product to the soil at a rate of 1 lb. per 100 sq. ft. (1 lb. per acre). Sprinkle over the soil uniformly in areas where these insects are seen or are known to exist. Reapply after it is consumed or as needed for continued control. Apply approximately 30 lbs. per acre.

Apply in late afternoon or early morning when insects have been seen or are known to exist.

MOLE CRICKETS in lawns, golf courses and ornamental turf - For control of Mole Crickets apply this granular product at the rate of 30 lbs. per acre or approximately 1 lb. per 100 sq. ft. spread evenly over the area using a mechanical spreader. Apply in late afternoon or moist soil in areas known to be infested with these insects. Repeat application is about one week for best control.

THE BEST TIME TO CONTROL THE SOUTHERN MOLE CRICKET

Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
	2/	1/	1/	1/	1/	1/	1/				

NOTE - #2 indicates the best time to control the Adult Mole Crickets.
#1 indicates the laying time during the summer