

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

JUL 20 2006

Thursday, July 20, 2006

Ms. Lisa Boynton
NCH Corp.
2727 Chemsearch Blvd.
Irving, TX 75062Subject: Submission of an amendment to modify existing label to add new uses and label claims.
EPA Registration Number: 1769-373
Product Name: Danco Concentrate
EPA Receipt Date: February 24, 2006

Dear Ms. Boynton;

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted prior to releasing the product for shipment.

1. Page 1 of 8:

- a. The product name on file must be identified as "Danco Concentrate" only. The "for Automated Misting Systems" should be placed in brackets, and identified as a footnote with "Optional Text", or identified as an alternate brand name.
- b. Revise "(for bulk containers)" to 55 gallons, or delete. Bulk Containers requires specific labeling language that is not presented on this label.
- c. On the right hand side of the label opposite the EPA Reg. No. 1769-373, insert an Establishment Number.¹

2. Page 2 of 8:

- a. For containers greater than the Environmental Hazards section revise the entire sections as follows:
 "[for containers less than 5 gallons]
 This product is toxic to fish. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.
 [for containers greater than 5 gallons]
 Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the

¹ LOCATION OF ESTABLISHMENT NUMBER. The establishment number must appear on either the product label or on the immediate container of the product. If the EPA Establishment number cannot be seen or clearly read through the wrapping or packaging of the pesticide container, it must also appear on the outside container or wrapper of the product [40 CFR 156.10 (f)].

permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.”²

3. Page 3 of 8:

- a. Within the Crawling Insects section, delete the language that is marked with a “strikeout”

~~CRAWLING INSECTS (such as Ants, Booklice, Carpet Beetles, Centipedes, Clothes Moths, Clover Mites, Cockroaches, Crickets, Dermestid Beetles, Earwigs, Firebrats, Fleas, Millipedes, Scorpions, Silverfish, Sowbugs and Spiders): Apply as general surface spray, crack and crevice treatment or void injection. Use the 0.25% or 0.50% rate.~~
~~GENERAL SURFACE SPRAY: Apply as a coarse, wet spray on walls, floors and other surfaces. Treat food storage areas, handling areas and unloading and processing areas.~~
~~CRACK AND CREVICE TREATMENT AND VOID INJECTION: Apply to cracks and crevices or structural voids in these places as well as beneath sinks and inside and around the base of food conveying, processing and handling equipment.~~

This text is already expressed in the methods of application at the beginning of the page. In addition, areas under sinks, around bases of conveying, processing and handling equipment are not considered void injection areas.

- b. At the end of the “BED BUGS, LICE AND TICKS:” section, insert the following text. “Allow mattresses and springs to thoroughly dry prior to reuse.”

4. Page 4 of 8:

- a. Within the subsection entitled “Mosquitoes” strike the following sentence:
~~Treatment of vegetation in water from land is also permitted by allowing wind to move product into vegetation, but application of this product directly to any body of water is prohibited.~~
- b. Remove the entire section “~~TRUCK MOUNTED EQUIPMENT: When using truck mounted equipment, spray at a dosage rate of fifteen (15) oz per minutes from a location up wind of the areas where insects are to be controlled. Keep speed of vehicle to approximately five (5) MPH. Figure application travel path for a convergence of a three hundred (300) foot swath. Repeat application as required for control.~~”

5. Page 5 of 8:

- a. Revise the Plant Safety Notice section as follows: “~~PLANT SAFETY NOTICE: PLANT PHYTOXICITY: This product has demonstrated excellent plant safety; however,~~ Aall varieties of the plants listed have not been tested for phytotoxicity. When treating large numbers of plants of a single variety, treat a few plants and observe for phytotoxicity prior to full scale application. Direct treatment to some open blooms may cause petal drop.”
- b. Remove the “but not limited to from the header “FOR USE INDOORS AND OUTDOORS ON, ~~BUT NOT LIMITED TO:~~”


² Further, PR Notice 95-1 exempts certain end-use products [i.e., products in containers of less than 5 gallons (liquid), less than 50 pounds (solid, dry weight) and in aerosol containers of any size] from bearing effluent discharge statements specified by PR Notice 93-10. PR Notice 93-10 still applies to the following kinds of pesticide products that may result in discharges to the waters of the United States or to municipal sewer systems, including but not limited to: (A) all technical grade and manufacturing use products; and (B) end-use products packaged in containers equal to or greater than 5 gallons (liquid) or 50 pounds (solid, dry weight), and registered for industrial preservative, water treatment, other industrial processing uses (such as cooling tower water systems, pulp and paper mill water systems, secondary oil recovery injection water systems, food processing operations, leather tanning, wood protection and textile treatment) and commercial and institutional uses (including, but not limited to, hospitals, hotels/motels, office buildings and prisons)

3/11

- c. Convert the statement "**Do not treat vegetables.**" to bold text.
- d. Within the Use Site listing, delete the following use site: "Stored food in multi-walled paper or cloth bags."

If you have any questions, you may contact Richard J. Gebken, at (703) 305-6701.

Sincerely,



Richard Gebken
Product Manager 10
Insecticide Branch
Registration Division (7505C)

Enclosure

4/11

**DANCO CONCENTRATE for Automated Misting Systems
PYRETHRIN CONCENTRATE**

KILLS: Angoumois Grain Moths, Ants, Boxelder Bugs, Cadelles, Cheese Mites, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Earwigs, Fruit Flies, Gnats, Granary Weevils, House Flies, Indian Meal Moths, Mosquitoes, Rice Weevils, Spiders, Tobacco Moths and Wasps.

FOR USE IN AND AROUND: Apartments, Cafeterias, Dormitories, Food Service and Preparation Areas, Grocery and Convenience Stores, Hotels, Health Care Facilities, Motels, Office Buildings, Restaurants, Schools and Universities.

ACTIVE INGREDIENTS

Pyrethrins	3.0%
Piperonyl Butoxide, Technical*	6.0%
n-Octyl Bicycloheptene Dicarboximide	10.0%
OTHER INGREDIENTS:	<u>81.0%</u>
TOTAL:	100.0%

*Equivalent to 4.8% (butylcarbityl)(6-propylpiperonyl) ether and 1.2% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

(See elsewhere for additional precautionary statements.)

ACCEPTED
with COMMENTS
on EPA Letter Dated

JUL 20 2006

NET CONTENTS: _____ (for bulk containers)
24 fl oz/708 ml
32 fl oz/946 ml

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, the pesticide
registered under EPA Reg. No.

1769-373

EPA Reg. No. 1769-373

EPA Est. No. 4140 -							
CA-1	IN-1	NP-1	PC-1	TX-1	TX-2	TX-4	CAN-1

NCH Corporation
2727 Chemsearch Blvd.
Irving, TX 75062
©2006 NCH Corporation

TO REORDER,
Call TOLL-FREE: 1-800-527-9919
FAX: 1-800-456-0634

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

For medical or environmental emergencies, call CHEMTREC® at 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

GENERAL RESTRICTIONS

Do not apply directly to food. Do not apply as a space or ULV spray while food processing is underway. Cover or remove any food in the treatment area before application. Cover all food processing surfaces and equipment or wash with an effective cleaning compound followed by a potable water rinse.

Food processing operations do not have to be stopped while applying as a wet spray with care and in accordance with directions and cautions above in establishments which do NOT operate under Federal meat, poultry, shell egg grading and egg products inspection programs.

In the home, cover all food processing surfaces, dishes and utensils in the treatment area or thoroughly wash before use.

Remove pets, birds, and cover fish aquariums and turn off air pump motor before spraying. In health care facilities or schools, remove patients or students prior to treatment. Thoroughly ventilate treated area for two hours before reoccupying treated areas.

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

DILUTION RATES

Use undiluted in mechanical sprayers which produce particles of aerosol size (80% of droplets less than 30 microns), or diluted in pump sprayers or automated misting systems. Use the 0.125% rate for flying insects only. Use the 0.25% rate for maintenance. Use the 0.5% rate for clean outs or high insect infestations. Use the 0.05% rate only for automated misting or fogging applications outdoors.

Pyrethrin % Desired	Amount of Concentrate per Gallon of Water*
0.050%	2 fl oz
0.125%	5 1/3 fl oz
0.250%	10 2/3 fl oz
0.500%	21 1/3 fl oz
1.000%	42 2/3 fl oz

* Do not dilute with spray oil.

NOTE: 1/3 fl oz = 2 tsp and 1/2 fl oz = 1 tbsp

6/11

APPLICATION INSTRUCTIONS

SPACE SPRAY (INDOORS): Apply using a mechanical aerosol generator which creates particles of aerosol size (80% of droplets less than 30 microns). Prior to application, close all doors and windows, shut off all ventilating equipment and extinguish all flames in the area to be treated. Do not remain in treated areas after application. Leave room(s) closed for at least one (1) hour. Ventilate thoroughly before re-entry.

CRACK AND CREVICE TREATMENT: Use diluted or undiluted. Inject DANCO CONCENTRATE into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. Apply in such a manner to limit dripping and run-off on structural surfaces.

VOID INJECTION: Use diluted or undiluted. Use to flush and kill insects harboring in closed or inaccessible voids. Calculate the cubic area of the void and treat at the rate of five to ten seconds per three cubic feet. Several holes may be required in long-running voids. Apply in such a manner to limit dripping and run-off on structural surfaces.

SPOT TREATMENT AND GENERAL SURFACE SPRAY: Use diluted or undiluted. Hold spray nozzle 18 to 24 inches from areas to be treated. Treat surfaces until moist, but in such a manner to limit dripping and run-off.

INDOOR TREATMENTS

FLYING INSECTS (such as Bees, Cluster Flies, Flies, Gnats, Hornets, Mosquitoes, Moths, Small Flying Moths, Wasps, Webbing Clothes Moths and Yellow Jackets): Apply as a space spray or general surface spray. **SPACE SPRAY:** apply undiluted at the rate of 0.5 fl oz of concentrate per 1,000 cu ft of space. Point nozzle upward and direct spray mist to all parts of the room. **GENERAL SURFACE SPRAY:** use the 0.125% or 0.25% rate and direct spray toward upper corners of room especially around windows and other light sources which attract these insects. Do not remain in treated area(s) after application. Ventilate thoroughly before re-entry.

CRAWLING INSECTS (such as Ants, Booklice, Carpet Beetles, Centipedes, Clothes Moths, Clover Mites, Cockroaches, Crickets, Dermestid Beetles, Earwigs, Firebrats, Fleas, Millipedes, Scorpions, Silverfish, Sowbugs and Spiders): Apply as general surface spray, crack and crevice treatment or void injection. Use the 0.25% or 0.50% rate. **GENERAL SURFACE SPRAY:** Apply as a coarse, wet spray on walls, floors and other surfaces. Treat food storage areas, handling areas and unloading and processing areas. **CRACK AND CREVICE TREATMENT AND VOID INJECTION:** Apply to cracks and crevices or structural voids in these places as well as beneath sinks and inside and around the base of food conveying, processing and handling equipment.

STORED PRODUCT PESTS (such as Angoumois Grain Moths, Caddies, Carpet Beetles, Chocolate Moths, Cigarette Beetles, Dermestid Beetles, Dried Fruit Beetles, Drug Store Beetles, Flour Beetles, Grain Mites, Granary Weevils, Indian Meal Moths, Meal Worms, Mediterranean Flour Moths, Psocids, Ref Flour Beetles, Saw-toothed Grain Beetles, Spider Beetles, Trogaderma and Weevils): Apply as a space spray or general surface spray. Clean all food preparation areas thoroughly after treatment and before food is exposed. Use only in areas where food is not exposed. **SPACE SPRAY:** apply undiluted at the rate of 0.5 fl oz of concentrate per 1,000 cu ft of space. Direct treatment toward upper corners of room and ceiling. **GENERAL SURFACE SPRAY:** use the 0.25% or 0.50% rate. Direct output into corners and under pallets, hitting as many insects as possible.

BED BUGS, LICE AND TICKS: Apply as a space spray, spot treatment or crack and crevice treatment. First, thoroughly clean air mattresses and springs. **SPACE SPRAY:** apply undiluted at the rate of 0.5 fl oz of concentrate per 1,000 cu ft of space. **CRACK AND CREVICE TREATMENT OR VOID INJECTION:** use the 0.25% or 0.50% rate and apply to cracks and crevices or structural voids around baseboards, floorboards, bedboards and walls. **SPOT TREATMENT:** Take bed apart. Treat joints and channels if hollow, such as square or round tubing. See that the interior framework is treated. Treat mattresses, especially tufts, folds and edges. Also treat other areas where bed bugs may be hiding.

7/11

FLEAS: Apply as a GENERAL SURFACE SPRAY undiluted or using the 0.25% or 0.50% rates. Direct treatment onto pet resting areas and floor around these areas. Treat pet bedding the same way. Also treat entire inside surface of dog houses. Do not allow pets to contact treated areas until spray has dried. Remove or cover animal feed before spraying. To control the source of flea infestations, pets inhabiting the treated premises should be treated with a product registered for application to pets.

DARKLING BEETLES: After emptying poultry houses, apply as a space spray or general surface spray. Treat litter before cleaning. Keep closed for one hour, remove litter and retreat all surfaces. Ventilate for 24 hours before replacing poultry. **SPACE SPRAY:** apply undiluted at the rate of 0.25 fl oz of concentrate per 1,000 cu ft of space. **GENERAL SURFACE SPRAY:** use the 0.25% or 0.50% rate.

INDOOR AND OUTDOOR TREATMENTS

BOXELDER BUGS, CARPENTER ANTS, CLUSTER FLIES AND ELMLEAF BEETLES: Apply as a CRACK AND CREVICE TREATMENT or as a void injection to voids and other areas where these insects are found. Use undiluted or at the 0.25% or 0.50% rates.

FLIES, GNATS, HORN FLIES, HOUSE FLIES, STABLE FLIES (AROUND LIVESTOCK): For fast knockdown of flying insects, apply as a SPACE SPRAY undiluted or using the 0.25% or 0.50% rates. Direct fog upward in all directions. Ventilate the area after treatment is completed. Remove or cover animal feed before spraying. Do not apply to lactating or food animals. Do not contaminate milk, food or drinking water.

OUTDOOR TREATMENTS

MOSQUITOES, FLIES, AND GNATS: Dilute to 0.050% or 0.125% when applying with mechanical, electrical, fogging or automated misting equipment, or when applying as a GENERAL SURFACE SPRAY. To control these pests in open areas near homes, buildings and in campgrounds, treat shrubbery and vegetation and other areas where these pests may rest. Structural surfaces, shrubbery, and vegetation around stagnant pools, marshy areas, ponds, and shorelines may be treated. Treatment of vegetation in water from land is also permitted by allowing wind to move product into vegetation, but application of this product directly to any body of water is prohibited.

AUTOMATED MISTING SYSTEMS: Automatic spraying systems must not be programmed to release pesticide where people, pets, food, or feed are directly exposed. A legible copy of the label must be securely attached to the equipment pesticide holding container or on the outside of the equipment timer.

TRUCK MOUNTED EQUIPMENT: When using truck mounted equipment, spray at a dosage rate of fifteen (15) oz per minutes from a location up-wind of the areas where insects are to be controlled. Keep speed of vehicle to approximately five (5) MPH. Figure application travel path for a convergence of a three hundred (300) foot swath. Repeat application as required for control.

HORNETS, WASPS AND YELLOW JACKETS: Apply as a GENERAL SURFACE SPRAY using the 0.25% or 0.50% rate. Application should be made in late evening when insects are at rest. Spray liberally into hiding and nesting places, contacting as many insects as possible.

HORTICULTURE USE

FOR USE IN: Hobby Greenhouses, Interiorscapes and Landscapes.

KILLS: Aphids, Armyworms, Bees, Beetles, Bugs (Boxelder, Lace and Plant), Caterpillars (Cabbage Worms), Rose Slugs, Hornworms), Centipedes, Clover Mites, Crickets, Earwigs, Firebrats, Flies, Fruitworm, Fungus Gnats, Fungus Gnat Larvae, Grasshoppers, Leafhoppers, Leafminers, Leaf Rollers, Loopers, Mealybugs, Millipedes, Mosquitoes, Psylia, Rose Aphids, Scale Insects, Scorpions, Sowbugs, Spiders, Spider Mites, Thrips, Wasps, Whiteflies and Yellow Jackets.

APPLICATION INSTRUCTIONS

This product is designed for diluted use (0.05% - 0.25%) in sprayers. DO NOT DILUTE PRODUCT WITH SPRAY OIL. Treat upper and lower foliage surfaces of plants uniformly.

SPRAY DILUTION SHART

Pyrethrin % Desired	Amount of Concentrate per Gallon of Water*
0.050%	2 fl oz
0.125%	5 1/3 fl oz
0.250%	10 2/3 fl oz

* Do not dilute with spray oil.

PLANT SAFETY NOTICE: This product has demonstrated excellent plant safety; however, all varieties of the plants listed have not been tested. When treating large numbers of plants of a single variety, treat a few plants and observe for phytotoxicity prior to full scale application.

Direct treatment to some open blooms may cause petal drop.

FOR USE INDOORS AND OUTDOORS ON, BUT NOT LIMITED TO:

Bedding plants (such as): impatiens, marigolds, petunias, geraniums, and garden mums.

Cut Flowers (such as): roses, chrysanthemums, carnations, snapdragons, and orchids.

Flowering hanging baskets (such as): fuchsia, lantana, New Guinea impatiens, ivy, and geraniums.

Foliage (such as): Boston fern, dracaena, ficus, schefflera, and philodendron.

Potted Flowering plants (such as): poinsettia, chrysanthemums, florist's azaleas, lilies, African violets, geraniums, hibiscus, kalanchoes, and begonias.

Ornamentals (such as): azalea, barberry, cotoneaster, euonymus, holly, juniper, oak, pine, rhododendron, roses, spiraea, spruce, viburnum and yew.

Do not treat vegetables.

In addition to sites listed on the front panel, DANCO CONCENTRATE is acceptable for use in the following sites:

INDOORS

- | | | |
|-------------------------------------|------------------------------|----------------------------|
| Animal areas and quarters | Food handling establishments | Milking parlors |
| Apartments | Food processing plants | Mills |
| Auditoriums | Food storage areas | Missions |
| Bakeries | Freight containers | Mobile homes |
| Barns | Fruit packing sheds | Motels |
| Beverage plants | Garbage cans | Museums |
| Boats | Garbage compactors | Mushroom houses |
| Bottling plants | Goat houses | Mushroom processing plants |
| Brandy storage warehouses | Grain elevators | Nursing homes |
| Buildings | Grain handling equipment | Office buildings |
| Buses | Grain mills | Passenger rail cars |
| Cabins | Granaries | Peanut processing plants |
| Calf hutches | Hobby greenhouses | Peanut storage warehouses |
| Canneries | Hog operations | Playgrounds |
| Cattle barns | Homes | Poultry houses |
| Churches | Horse barns | Poultry operations |
| Closets | Horse stables | Poultry processing plants |
| Commercial and industrial buildings | Hospices | Prisons |
| Communication centers | Hospitals | Public buildings |
| Condominiums | Hotels | Rabbit houses |
| Correctional facilities | Industrial installations | Rabbit processing plants |
| Dairies | Institutions | Railroad cars |
| Dairy and livestock barns | Interiorscapes | Recreational vehicles |
| Drains and sewers | Jails | Research animal areas |
| Dried fruit processing plants | Kennels | Restaurants |
| Dried fruit warehouses | Laboratories | Restrooms |
| Drugstores | Libraries | Rice mills |
| Dumpsters | Livestock barns | Schools |
| Dwellings | Loafing sheds | Shade houses |
| Egg processing plants | Locker rooms | Shipholds |
| Factories | Mausoleums | Ships |
| Flour mills | Meat packing plants | Stable bins |
| | Milk rooms | Stables |

paper or cloth bags
Store seed warehouses
Stores
Supermarkets
Swine houses
Textile mills and warehouses
Theaters

Tobacco warehouses
Townhouses
Trains
Transportation equipment
Truck trailers
Trucks
Utilities

Voids (attics, walls, ceilings)
Warehouses
Wine storage warehouses
Wineries
Zoos

OUTDOORS

Campgrounds
Corrals
Drive-in restaurants
Drive-in theaters
Dumpsters
Feedlots
Garbage dumps
Golf courses
Junkyards
Kennels

Landscapes
Lath houses
Manure piles
Parks
Plantscapes
Recreational areas
Sewers
Swineyards
Zoos

DANCO CONCENTRATE is also effective against the following insects when used according to label directions:

FLYING INSECTS

Aphids
Barn Flies
Bed Bugs
Bees
Black Flies
Blow Flies
Bottle Flies
Cheese Skippers
Cluster Flies
Deer Flies
Face Flies
Flies
Fruit Flies
Fungus Gnats
Gnats

Horn Flies
Hornets
Horse Flies
House Flies
Mosquitoes
Mud Daubers
Mushroom Flies
Skipper Flies
Stable Flies
Thrips
Vinegar Flies
Wasps
Whiteflies
Yellow Jackets

GRAIN BORDERS

Larger Grain Borers

Lesser Grain Borers

CRAWLING INSECTS

Ants
Bed Bugs
Booklice
Boxelder Bugs
Cabbage Loopers
Cabbage Worms
Carpenter Ants
Centipedes
Chiggers
Chrysanthemum Leaf Miners

Cockroaches
Crickets
Earwigs
Fire Ants
Firebrats
Grasshoppers
Mealy Bugs
Millipedes
Palmetto Bugs
Pill Bugs

10/11

Psocids
Scorpions
Silverfish

FLEAS AND TICKS

Brown Dog Ticks
Fleas
Sheep Ticks

SPIDERS AND MITES

Cheese Mites
Clover Mites
Grain Mites

LICE

Lice

WORMS

Dark Mealworms
Lesser Mealworms

BEETLES

American Black Flour Beetles
Black Carpet Beetles
Black Fungus Beetles
Broadhorned Flour Beetles
Brown Spider Beetles
Cadelles
Carpet Beetles
Catorama Beetles
Cereal Beetles
Cigarette Beetles
Confused Flour Beetles
Corn Sap Beetles
Darkling Beetles
Depressed Flour Beetles
Dermestid Beetles
Dried Fruit Beetles
Drugstore Beetles
Elmleaf Beetles
Flat Grain Beetles
Flour Beetles
Foreign Grain Beetles
Fungus Beetles

Sowbugs
Spring Tails
Vegetable Leaf Miners

Spinosa Ear Ticks
Ticks

Red Spider Mites
Spider Mites
Spiders

Mounting stages of yellow lice and sucking blue lice

Mealworms
Yellow Mealworms

Ground Beetles
Hairy Fungus Beetles
Hairy Spider Beetles
Hide Beetles
Khapra Beetles
Large Black Flour Beetles
Longheaded Flour Beetles
Merchants Grain Beetles
Mexican Grain Beetles
Murmidius Beetles
Red Flour Beetles
Red Horned Grain Beetles
Rusty Grain Beetles
Saw-Toothed Grain Beetles
Slenderhorned Flour Beetles
Smalleyed Flour Beetles
Spider Beetles
Squarenecked Grain Beetles
Trogoderma Beetles
Two Banded Beetles
White Marked Spider Beetles

11/11

WEEVILS

Bean Weevils
Broadnosed Grain Weevils
Coffee Bean Weevils

Granary Weevils
Maize Weevils
Rice Weevils

MOTHS

Almond Moths
Angoumois Grain Moths
Chocolate Moths
Clothes Moths
Cocoa Bean Moths
European Grain Moths
Indian Meal Moths

Mediterranean Flour Moths
Moths
Pink Scavenger Caterpillars
Rice Moths
Small Flying Moths
Tobacco Moths
Webbing Clothes Moths

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container in secured dry storage area. Do not store above 100°F for extended periods of time. Storage below 32°F may cause freezing.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Empty container by using the product in accordance with the label directions. Do not reuse this container! Place empty container in trash or offer for recycling if available. If container is partly filled, call your local solid waste agency or call 1-800-CLEANUP for disposal instructions. NEVER PLACE UNUSED PRODUCT DOWN ANY INDOOR OR OUTDOOR DRAIN!

MADE IN THE U.S.A.

Label Code

oM602

