

CHLORDANE: When mixed with water this product forms a water suspended, residual type insecticide. The amount of water used will depend on available spraying equipment.

DIRECTIONS

LAWN INSECTS: For the control of ants, chiggers, crickets, grasshoppers, boxelder bugs and spiders, apply 3 to 4 pints per acre. For the control of chinch bugs, Japanese beetle grubs, white-fringed beetle larvae, white grubs and wireworms, apply 5 to 6 pints per acre.

For small areas add the following amount of Chlordane to the volume of water needed to cover the area.

CAUTION: Keep children and pets off treated areas until material is washed into the soil and the grass is thoroughly dry.

AMOUNT REQUIRED FOR NUMBER OF SQUARE FEET TO BE COVERED

Needed Per Acre	1,000 Sq. Ft.	5,000 Sq. Ft.	10,000 Sq. Ft.
2 pts. ✓	2 tbsp. ✓	10 tbsp. ✓	20 tbsp. ✓
4 pts. ✓	4 tbsp. ✓	20 tbsp. ✓	40 tbsp. ✓


EXAMPLE: you have a lawn with 4,000 sq. ft. and you find that you can cover it with 3 gallons of spray and the directions recommend 2 pints Chlordane per acre. You will then add 8 level tablespoons of Chlordane to 3 gallons of water and have the right amount of the right strength to spray your lawn.

HOUSEHOLD INSECTS: For spot application only. For the control of roaches or water bugs, silverfish, ants and spiders, mix 10 tablespoons Chlordane per gallon of water. For roaches spray thoroughly the infested areas such as cracks, crevices, along water pipes and darkened areas which serve as hiding places.

For silverfish spray infested areas of walls, floors, and shelves of linen and clothes closets, around water pipes, under and around lavatories and sinks and to the underside of book shelves.

When treating for ants and spiders special emphasis should be placed on the treatment of ant trails, and around door frames and window sills where pests enter premises. Thoroughly work spray into infested cracks and crevices.

CONTENTS ONE GALLON



18-1237



CHLORDANE

4 CONTAINS 4 LBS. TECHNICAL CHLORDANE PER GALLON 4

ACTIVE INGREDIENTS:

*Technical Chlordane	45%
Aliphatic Petroleum Solvent	51%
INERT INGREDIENTS	4%
	100%

*Equivalent to 27% octachloro-4, 7-methanotetrahydroindane, and 18% related compounds.

MIXES WITH WATER IN ALL PROPORTIONS

CAUTION

Keep out of reach of children
See side panel for additional precautionary labeling

E.P.A. Reg. No. 1990-178

DIRECTIONS

CRABGRASS CONTROL: Apply 3 pints per each 1,000 sq. is important. Chlordane must be applied before crabgrass before May 1. Do not apply to newly seeded lawns. After wash the Chlordane off grass and down onto the surface children or pets on the treated area until the Chlordane and the area has dried.

SUBTERRANEAN TERMITES — Directions for Non-Professional in a 1.0% water emulsion, 4½ qts. Chlordane in 50 gallon **Buildings Having Basements or Crawl Spaces:** Dig a 1 of the foundation walls. In brick or hollow block or concrete trench to, but not below, the footing. Then as the trench soil at the rate of 4 gallons per 10 linear feet for each 3 feet deep would require 12 gallons per 10 linear feet. The objective is to establish a treated soil barrier which will Other critical areas may also require treatment. It is suggested the State Extension Service for advice.

FRUIT INSECTS: Apples, peaches, pears, for the control of rose chafer and flea beetles. Use 2 pints per 100 gallons of plum curculio apply as a thorough cover spray beginning stage. Do not apply within 30 days of harvest.

CAUTION: Contact with skin can cause toxic symptoms. Avoid and skin contact. In case of spillage on skin, wash immediately with water. Avoid contamination of feed and foodstuffs. Do not or humans. Harmful if swallowed. Keep out of reach of to dairy or meat animals.

Do not use in edible products areas of food processing or other areas where food is commercially prepared or prepared serving areas while food is exposed.

This product is highly toxic to bees exposed to direct contact with crops. Protective information may be obtained from your Extension Service.

This product is toxic to fish, birds, and other wildlife. areas may be killed. Do not apply where runoff is likely when weather conditions favor drift from areas treated. Do by cleaning of equipment, or disposal of wastes.

NOTICE: Do not reuse container. Wash thoroughly with water in a safe place.

NOTE: DO NOT USE, POUR, SPILL OR STORE NEAR HEAT. **NON-WARRANTY:** Buyer assumes all risks of use, storage, or not in strict accordance with directions given herewith.

FARMLAND INDUSTRIES

This product forms a water sus-
amount of water used will depend

ONS

ants, chiggers, crickets, grass-
apply 3 to 4 pints per acre.
ese beetle grubs, white-fringed
worms, apply 5 to 6 pints per

unt of Chlordane to the volume

treated areas until material is
is thoroughly dry.

SQUARE FEET TO BE COVERED

5,000	10,000
Sq. Ft.	Sq. Ft.
10 tbsp.	20 tbsp.
20 tbsp.	40 tbsp.

0 sq. ft. and you find that you
y and the directions recommend
ll then add 8 level tablespoons
nd have the right amount of the

ion only. For the control of roaches
ers, mix 10 tablespoons Chlordane
ray thoroughly the infested areas
r pipes and darkened areas which

f walls, floors, and shelves of
water pipes, under and around
erside of book shelves.

pecial emphasis should be placed
around door frames and window.
Thoroughly work spray into in-

E GALLON


0-178

TED

1973

PESTICIDE
IDE ACT
GISTER-
SUBJECT
RS.

18-1237



CHLORDANE

4 CONTAINS 4 LBS. TECHNICAL CHLORDANE
PER GALLON 4

ACTIVE INGREDIENTS:

*Technical Chlordane	45%
Aliphatic Petroleum Solvent	51%
INERT INGREDIENTS	4%
	100%

*Equivalent to 27% octachloro-4, 7-methanotetrahydroindane,
and 18% related compounds.
MIXES WITH WATER IN ALL PROPORTIONS

CAUTION

Keep out of reach of children

See side panel for additional precautionary labeling

E.P.A. Reg. No. 1990-178

DIRECTIONS

CRABGRASS CONTROL: Apply 3 pints per each 1,000 sq. ft. of lawn area. Timing is important. Chlordane must be applied before crabgrass seed germinates, usually before May 1. Do not apply to newly seeded lawns. After application immediately wash the Chlordane off grass and down onto the surface of the soil. Do not allow children or pets on the treated area until the Chlordane has been watered down and the area has dried.

SUBTERRANEAN TERMITES — Directions for Non-Professional Use. Use Chlordane in a 1.0% water emulsion, 4½ qts. Chlordane in 50 gallons of water, as follows:
Buildings Having Basements or Crawl Spaces: Dig a trench along the outside of the foundation walls. In brick or hollow block or concrete foundations, dig a trench to, but not below, the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons per 10 linear feet for each foot of depth. A trench 3 feet deep would require 12 gallons per 10 linear feet.

The objective is to establish a treated soil barrier which will prevent termite entry. Other critical areas may also require treatment. It is suggested that you consult the State Extension Service for advice.

FRUIT INSECTS: Apples, peaches, pears, for the control of plum curculio, thrips, rose chafer, and flea beetles. Use 2 pints per 100 gallons of water. For the control of plum curculio apply as a thorough cover spray beginning with the shuck split stage. Do not apply within 30 days of harvest.

CAUTION: Contact with skin can cause toxic symptoms. Avoid breathing spray mist and skin contact. In case of spillage on skin, wash immediately with soap and water. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Harmful if swallowed. Keep out of reach of children. Do not apply to dairy or meat animals.

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

NOTICE: Do not reuse container. Wash thoroughly with water and detergent. Discard in a safe place.

NOTE: DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

NON-WARRANTY: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

FARMLAND INDUSTRIES, INC.

KATONAH, MISSOURI 64563