

USDA REG. NO. 1772-42

SEVIN* DUST

CONTROLS

JAPANESE BEETLE, MEXICAN BEAN BEETLE, FLEA BEETLE, LEAF-HOPPER, CUCUMBER BEETLE, SQUASH BUG, STINK BUG, COLORADO POTATO BEETLE, TOMATO FRUITWORM, HORNWORM, CORN EARWORM AND CERTAIN OTHER INSECTS, ON THE PLANTS NOTED IN THE TABLE ON THE REVERSE SIDE.

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-Methylcarbamate) 2.0%

INERT INGREDIENTS 98%

100%

*Trade Mark of Union Carbide Corporation

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

This product is harmful if inhaled or swallowed. Avoid breathing the dust or spray and avoid prolonged and repeated contact with the skin. Wash thoroughly after handling. Wear regular long-sleeved work clothing and change to clean clothing daily. Avoid contamination of food or feedstuffs. Do not apply the dust or spray to edible plants within 7 days before gathering. Do not feed the treated plants to animals. Do not reuse the empty container. Bury it with waste or burn it. Stay away from smoke and fumes. **NOTE TO PHYSICIANS:**

AMBUCRYL is a moderate, reversible cholinesterase inhibitor. Atropine is antidote. **THIS PRODUCT IS TOXIC TO BEES.** Any use of it around bees, or on the crop, or to the tassel region of corn, will seriously reduce bee populations. Bees are important pollinators and are essential to the production of many crops. Before using this product during such periods, consult your State Agricultural Extension Service.

Holder Corporation

HUNTINGTON, WEST VIRGINIA

25 LBS. NET

PLANT	PEST	DOSAGE	
		Per Acre	Per 1,000 Sq. Ft.
Bean	Mexican Bean Beetle	50 lbs.	1 lb. 2 1/4 ozs.
	Flea Beetle, Leafhopper	50 lbs.	1 lb. 2 1/4 ozs.
Cucumber	Cucumber Beetle,	50 lbs.	1 lb. 2 1/4 ozs.
	Pickleworm	50 lbs.	1 lb. 2 1/4 ozs.
Potato	Colorado Potato Beetle,	50 lbs.	1 lb. 2 1/4 ozs.
	Flea Beetle, Leafhopper	50 lbs.	1 lb. 2 1/4 ozs.
Summer Squash	Cucumber Beetle, Pickleworm,	50 lbs.	1 lb. 2 1/4 ozs.
	Squash Bug	50 lbs.	1 lb. 2 1/4 ozs.
Sweet Corn	Corn Earworm	75 lbs.	1 lb. 11 1/4 ozs.
Tomato	Tomato Fruitworm, Hornworm,	60 lbs.	1 lb. 6 ozs.
	Stink Bug	60 lbs.	1 lb. 6 ozs.
On the above, and on			
Ornamentals	Japanese Beetle	50 lbs.	1 lb. 2 1/4 ozs.

DIRECTIONS

The above dosages are recommended. At the first appearance of the insect or worm (except in the case of sweet corn) dust the plants lightly but thoroughly, covering both sides of the leaves. Repeat as often as necessary.

SWEET CORN: When the silks appear, dust the silk and ear, and repeat 3 or 4 times at 2-day or 3-day intervals. To prevent budworm damage, dust the whorls of the plant before the silks or tassels appear. **IF THIS MATERIAL IS TO BE USED AS A SPRAY,** mix thoroughly 6 ozs. to each gallon of water and follow the schedule for dusting.

