

**INGREDIENTS**

50 SEVIN furnished by Union Carbide Corp. (New York) designated by them as EC, and which contains a wetting and dispersing agent.

50 MC

**DIRECTIONS**

PLANT	PEST	Level table-spoonfull per gallon of water	Approx. dosage per 1,000 sq. ft.
Bean	Mex. Bean Beetle, Flea Beetle, Leafhopper .....	2	¾ ozs.
	Cucumber	Cucumber Beetle, Pickleworm ...	2
Potato	Colorado Pot. Beetle, Flea Beetle, Leafhopper .....	2	¾ ozs.
	Summer Squash	Cucumber Beetle, Pickleworm, Squash Bug .....	2
Sweet Corn	Corn Earworm .....	3	1 ¼ ozs.
Tomato	Tomato Fruitworm, Hornworm, Stink Bug .....	2 ½	¾ ozs.
	Ornamentals	Bagworm, Japanese Beetle, Tent Caterpillar .....	2

EPA Reg No. 1772-43

**SEVIN\* 50-W**

**CONTROLS**

MEXICAN BEAN BEETLE, JAPANESE BEETLE, FLEA BEETLE, LEAF HOPPER, CUCUMBER BEETLE, SQUASH BUG, STINK BUG, COLORADO POTATO BEETLE, TOMATO PICKLEWORM, HORNWORM, CORN EARWORM ON THE PLANTS NOTED IN THE TABLE ON BACK PANEL

ACTIVE INGREDIENT	Carbaryl (1-Naphthyl N-Methylcarbamate)	50%
INERT INGREDIENTS		50%
		100%

\*Trade Mark of Union Carbide Corporation


**CAUTION**

**KEEP OUT OF THE REACH OF CHILDREN**

This product is harmful if swallowed or inhaled. Avoid contact with the mucous membrane of the spray mist. Wear regular long-sleeved clothing and wash thoroughly after handling. Do not contaminate food or feed stuffs. Do not feed treated plants to animals. Do not permit children to play on or near treated plants. Note to Physician: Carbaryl is a cholinesterase inhibitor; atropine is antidotal.

This product is harmful to bees. Application to the tassels region of corn should be avoided during the flowering period, or to other plants when bees are active. This will severely reduce the bee population. Before spraying during such periods, contact your State Extension Service.

Use as a spray only. Follow the dilution and dosage recommended in the table. At the first appearance of the insect or worm (except in the case of sweet corn spray the plants lightly but thoroughly, covering both sides of the leaves. Repeat at weekly or 10-day intervals as needed. SWEET CORN: To control budworm damage, spray the silks as soon as they appear. To control earworm, when the silks first appear, spray the silk and ear thoroughly and repeat over 4 times at 2- or 3-day intervals.



**Holder Corporation**

HUNTINGTON, WEST VIRGINIA