

6% CHLORDANE DUST

CONTROLS ANTS, WIREWORMS, WHITE GRUBS, JAPANESE BEETLE GRUBS, On Plants Noted In Directions.

Active Ingredient:	
Technical Chlordane*	6%
Inert Ingredients	94%
	100%

*Equivalent to 3.6% Octachloro-4,7-Methanotetrahydroindane, and 2.4% related compounds.

DIRECTIONS

Beans, Sweet Corn, Cucumbers, Cantaloups, Potatoes, Peppers, Summer Squash, Tomatoes, and Onions: To control Wireworms, broadcast 70 to 140 lbs. per acre (2½ to 5 ozs. per 100 sq. ft.) on soil before planting and work into the top 6" with a discer or cultivator. Do not repeat application for 3 years.

Strawberries: To control Wireworms, broadcast 70 to 140 lbs. per acre (2½ to 5 ozs. per 100 sq. ft.) on soil before planting and work into top 6" with a discer or cultivator. Do not repeat application for 3 years.

Lawns and Turf not used for grazing: To control Ants, Japanese Beetle Grubs, White Grubs, Chinch Bugs, European Chafer, Army Worms, Cicada-Killer Cutworms: Broadcast 3 to 4 lbs. per 1000 sq. ft. (5 to 7 ozs. per 100 sq. ft.) when insects are first observed and wash into soil after application. Do not apply when ground is frozen.

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

Do not allow dust to drift when applying. Wear gloves and protective clothing. Avoid breathing the dust or prolonged contact with the skin. Wash thoroughly with soap and water after handling. Do not allow the material to contaminate food or feedstuffs. Keep children and pets off treated areas until material is watered into the soil and the grass is dry. Harmful if swallowed.

This product is poisonous to fish and wildlife. Do not contaminate lakes, streams, or ponds. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift of dust from areas treated. Do not reuse container. Dispose of it safely.

Holder Corporation

HUNTINGTON, WEST VIRGINIA