

[1772-71]

Net Weight

25 lb.



ACCEPTED
4-17-67
456-2

"7.5" Rotenone

DIRECTIONS

Apply direct from package with an efficient type of duster. For normal infestations of Mexican Bean Beetle, Cabbage Worms and Colorado Potato Bugs use 20-30 lbs. per acre or 1 lb. each 600 feet of row. Increase for very heavy infestations. Cover all parts of the plant including both sides of leaf thoroughly. Effective application is more beneficial than large quantities. Repeat every 7 to 10 days or as required. Avoid dusting before rain storms. For best results dust in early morning or late afternoon. To use as a spray add 2 tablespoons to a gallon of water or 1 lb. in 5 gallons of water and mix thoroughly.

DO NOT APPLY WITHIN ONE DAY OF HARVEST

CONTROLS

- | | |
|---------------------|---------------------|
| Mexican Bean Beetle | Tobacco Flea Beetle |
| Cabbage Worms | Colorado Potato Bug |
| Cucumber Beetles | Tomato Horn Worm |
| Flea Beetles | Turnip Aphids |

Seller makes no warranty express or implied concerning the use of this product. Buyer assumes all risk of use whether in accordance with directions or not.

ACTIVE INGREDIENTS:

Rotenone	0.75
Other Cube Resins:	1.65

INERT INGREDIENTS:

Talc or Clay (Insecticidal)	97.60
Total	100.00

CAUTION: Keep Out of Reach of Children
See Other Warnings on Back Panel.