

[1772-78] 2 PAGES

Net Weight

4 Pounds

ACCEPTED

JAN 10 1968

UNDER THE FEDERAL
FUNGICIDE AND RODE
FOR ECONOMIC BOND
ED UNDER NO. 456-47
TO ATTACHED COM



"100" Rotenone

U.S.D.A. Reg. No. 456-47

ACTIVE INGREDIENTS:

Rotenone	1.00%
Other Cube Resins	2.50%

INERT INGREDIENTS	96.50%
--------------------------------	--------

Total	100.00%
--------------------	---------

CAUTION: Keep Out of Reach of Children
See Other Warnings on Back Panel.

ACCEPTED

JAN 10 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 47 SUBJECT
TO ATTACKED COMMENTS

"100" ROTENONE

DIRECTIONS:

Use as a dust in an efficient euster. Start applications when insects or damage first appear, give emphasis on coverage of both sides of leaves. Repeat applications every 7 - 10 days or oftener for severe infestations. May be used as a spray by mixing 1 lb. in 5 gallons of water and mixing thoroughly.

CROP	INSECTS CONTROLLED	DOSAGE AND LIMITATIONS
<u>beans - (all types)</u>	Mexican bean beetles, Flea beetle, <u>Striped Cucumber Beetle</u>	30-50 lbs. per acre. Do not use within 1 day of harvest.
<u>tomatoes</u>	Tomato hornworm, Flea beetle	30-50 lbs. per acre. Do not use within 1 day of harvest.
<u>Cabbage</u>	Striped Cucumber beetle Cabbage looper	Use 30-50 lbs. per acre. Do not apply within 1 day of harvest.
<u>Tobacco</u>	Flea beetles	Use 30-50 lbs. per acre. See the limitations.

CAUTION:

Avoid prolonged or repeated breathing of dust. Keep out of reach of children.

Manufacturer generally warrants material to be effective for indicated uses. Failure of buyer to properly use, handle and store said material voids all warranties.