Herculite V

Concentrated Disinfectant Detergent

Pseudomonacidal, Staphylocidal, Fungicidal

USE DIRECTIONS

Use at 2 ounces per gallon. This product will be effective against Pseudomonas aeruginosa PDR-10 at this level. This has been substantiated by the AOAC Use Dilution Confirmation Test. At this level, the product is fungicidal against pathogenic fungi.

Herculite V is a powerful disinfectant effective against the following organisms using AOAC use dilution test methods. Use dilution 2 oz. per gallon of water against Staphylococcus Aureus, Pseudomonas Aeruginosa, Salmonella Choleraesuis, Trichophyton Interdigitale.

For best results, use Herculite V with a cloth or mop and apply to walls, floors and other hard surfaces.

If used for sanitization of food equipment or food contact items, limit the level of active quaternary to 200 ppm. (2.8 ounces per 5 gallons). NO RINSE REQUIRED.

For heavy soil or organic matter, a precleaning step is recommended.

NET CONTENTS ____ GALLONS EPA Reg. No. 1778-49

FOR HOSPITALS, NURSING HOMES AND SCHOOLS

DANGER: Keep Out of the Reach of Children. See right column for antidote and other precautions.

Rinse empty container thoroughly with water and discard.

Manufactured by

WESTERN CHEMICAL COMPANY General Office, St. Joseph, Mo. 64501 VIRUCIDAL against HERPES SIMPLEX, ADENOVIRUS TYPE 2, INFLUENZA A₂ and VACCINIA on inanimate environmental surfaces.

Herculite V is a cleaner, disinfectant and sanitizer designed for institutional use such as hospitals, schools, nursing homes, where housekeeping is of prime importance in controlling the hazard of cross infection.

Herculite V can provide the assurance you need in this area, when used as directed. For use on walls, floors, sink tops, garbage pails, rest rooms, bed pans, operating rooms. DANGER: Keep Out of the Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink PROMPTLY a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities f water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

