

INSECTICIDE, MALATHION 1% DUSTING POWDER

ACTIVE INGREDIENTS: PERCENT BY WEIGHT

MALATHION
(0,0-dimethyl phosphorodithioate
of diethyl mercaptosuccinate) 1%

INACTIVE INGREDIENT:
PYROPHYLLITE 99%

CAUTION: KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS
USDA REG. NO. 1783-59

AVOID PROLONGED BREATHING OF DUST.

25 LB. NET

C 3/69 LOT 554

WT 30 CU 1.1

TRIO CHEMICAL WORKS, INC.
BROOKLYN, N. Y.

FOR MILITARY USE ONLY

STORAGE INSTRUCTIONS

~~Store in a cool, dry place when possible.~~ Never exceed 90°F. temperature. The average effective period of this item is 12 months from date of manufacture. If used after this period, malathion content should not be below 0.75 percent.

CAUTION!

For external use only. Keep away from food and mess gear. 20 ft. or face mask should be used if powder is dispensed into air gun or blower and one hour allowed for dust to settle before entering. The following additional precautions should be observed:

1. When dusting by hand or machine, do not apply more than 1/2 ounce to each person.
2. Do not redust for fourteen full days after previous dusting.
3. Individual 3-ounce polyethylene bottles of Malathion Powder, Dusting will not be used by personnel who have been dusted by hand or machine for at least fourteen full days after such dusting.
4. If possible, troops should bathe before being dusted.
5. Do not dust individuals who have insect repellent or grease on the skin.

INSTRUCTIONS FOR USE

FOR BODY LICE:

1. Clothing off person: Turn clothing inside out and dust 1/2 ounce over entire inner surface, rubbing powder well into all seams and creases. Repeat at weekly intervals until infestation is controlled.
2. Clothing on person: Use powder or hand duster. If hand duster, fill 3/4 full and adjust to deliver heavy cloud of powder at 12 inch stroke. (a). Dust inside of hat and replace on head. (b). Enter at side, shoulder height. Insert delivery tube slowly up each arm between skin and inner garments. Blow powder into armpit and shoulder, also between individual layers of clothing. (c). Insert tube at back of neck and dust back and neckband. (d). Insert tube at front at neck and dust toward armpits, both sides of body, chest and abdomen. (e). Loosen trousers, insert tube between skin and inner garments, and powder generously crotch and pubic area, also drawers, especially waist and side seams. (f). Insert tube between leg of trousers next to skin and powder buttocks and rear of crotch. (g). Dust in socks.

FOR HEAD AND PUBIC LICE: Dust into infested hairy portions of body and work down to the base of the hair with finger tips. Repeat when reinfestation occurs.

FOR FLEAS: For flea control in buildings, apply to cracks, crevices and rodent runways. Apply to floor areas as a thin, uniform coating of powder at the rate of 1/4 to 1/2 pound per 1,000 square feet. For infested grounds, apply the powder at the rate of 1 pound per 1,000 square feet or 40 pounds per acre.

ACCEPTED

MAY 9 1969

UNDER THE FEDERAL INSECTICIDE AND RODENTICIDE ACT OR ECONOMIC POISON REGISTERED UNDER NO. 1783-59 SUBJECT TO ATTACHED COMMENTS.

STORAGE INSTRUCTIONS

~~Store in a cool, dry place when possible.~~ Never exceed 90°F. storage temperature. The average effective period of this item is 12 months from date of manufacture. If used after this period, malathion content should not be below 0.75 percent.

CAUTION!

For external use only. Keep away from food and mess gear. Respirator or face mask should be used if powder is dispensed into air with dust gun or blower and one hour allowed for dust to settle before reentering. The following additional precautions should be observed:

1. When dusting by hand or machine, do not apply more than one ounce to each person.
2. Do not redust for fourteen full days after a previous dusting.
3. Individual 3-ounce polyethylene bottles of Malathion Powder, Dusting will not be used by personnel who have been dusted by hand or machine for at least fourteen full days after such dusting.
4. If possible, troops should bathe before being dusted.
5. Do not dust individuals who have insect repellents, oil or grease on the skin.

INSTRUCTIONS FOR USE

FOR BODY LICE:

1. Clothing off person: Turn clothing inside out and dust lightly over entire inner surface, rubbing powder well into all seams. Repeat at weekly intervals until infestation is controlled.
2. Clothing on person: Use powder or hand duster. If hand duster fill 3/4 full and adjust to deliver heavy cloud of powder at each stroke. (a). Dust inside of hat and replace on head. (b). Extend arms at side, shoulder height. Insert delivery tube slowly up each sleeve between skin and inner garments. Blow powder into armpit and about shoulder, also between individual layers of clothing. (c). Insert tube at back of neck and dust back and neckband. (d). Insert tube in front at neck and dust toward armpits, both sides of body, chest, and abdomen. (e). Loosen trousers, insert tube between skin and innermost garments, and powder generously crotch and pubic area, also underdrawers, especially waist and side seams. (f). Insert tube down rear of trousers next to skin and powder buttocks and rear of crotch. (g). Dust in socks.

FOR HEAD AND PUBIC LICE: Dust into infested hairy portions of the body and work down to the base of the hair with finger tips. Repeat when reinfestation occurs.

FOR FLEAS: For flea control in buildings, apply to cracks, crevices, and rodent runways. Apply to floor areas as a thin, uniform film of powder at the rate of 1/4 to 1/2 pound per 1,000 square feet. For treating infested grounds, apply the powder at the rate of 1 pound per 1,000 square feet or 40 pounds per acre.

ON

ACCEPTED

19 1969

FEDERAL INSECTICIDE
AND RODENTICIDE ACT
TOXIC POISON REGISTER
01783-48 SUBJECT
D COMMENTS.

C.

Y