



1791-28

**Cold Disinfection**

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1 1/2 ounce **RINSOL** per gallon of water for 10 minutes.

**Restaurant and Bar Rinse**

Dishes, glassware, silverware, cooking utensils: wash with soap or synthetic detergent; rinse thoroughly and immerse in a sanitizing solution containing 1 ounce **RINSOL** to 4 gallons of water. No Terminal Potable Water Rinse Required.

**FOR ROOM FOGGING**

**DIRECTIONS:**

To minimize the danger of cross-contamination from environmental surfaces apply solution of **RINSOL** as a fog as described below, before applying standard routine terminal cleaning and disinfecting practices. 1. Remove all human, animal, and plant life from room. 2. Open closet doors and drawers. 3. Set up 34" revolving platform in center of room. 4. Mount Jet FOGMASTER TRI-JET MODEL 6208, or other fogging device delivering equivalent spray at 3 rpm. 5. Fill sprayer reservoir with 1.5 ounces of **RINSOL** to 1 gallon of water to produce 1150 ppm active ingredients. 6. Set sprayer mechanism to deliver 1 gallon of solution. 7. Fog for 15 minutes for an average 2100 cu. ft. room. For different room sizes, vary spray time proportionately to ensure complete wetting of exposed surfaces. 8. Wait 2 hours before entering room after treatment. 9. Rinse fogging equipment thoroughly with clear water following use.

**DANGER**

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

**FIRST AID**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

**NOTE TO PHYSICIAN:**

Probably mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.



**DISINFECTANT** **RINSOL** **SANITIZER**

**ACTIVE INGREDIENTS:**

- n-Alkyl (60% C<sub>14</sub>, 30% C<sub>16</sub>, 5% C<sub>12</sub>, 5% C<sub>18</sub>) dimethyl benzyl ammonium chlorides.....
- n-Alkyl (68% C<sub>12</sub>, 32% C<sub>14</sub>) dimethyl ethylbenzyl ammonium chlorides .....
- Ethyl Alcohol .....
- INERT INGREDIENTS.....

**RINSOL** is a concentrate and should be diluted before use.

NET CONTENTS \_\_\_\_\_ EPA Reg. No. \_\_\_\_\_  
EPA Est. No. \_\_\_\_\_

**DANGER**

**KEEP OUT OF REACH OF CHILDREN**

See other precautionary and first aid statements on left



28

tools and dental equipment,  
RINSOL per gallon of water for

or synthetic detergent; rinse  
RINSOL to 4 gallons of water.

tal surfaces apply solution of  
routine terminal cleaning and  
e from room. 2. Open closet  
ter of room. 4. Mount Jet  
ing equivalent spray at 3 rpm.  
of water to produce 1150 ppm  
of solution. 7. Fog for 15  
vary spray time proportionately  
rs before entering room after  
er following use.

d skin damage. Do not get in  
dling concentrate. Harmful if

water for at least 15 minutes.  
ing before reuse. If swallowed,  
drink large quantities of water.

tric lavage. Measures against  
needed.



# DISINFECTANT RINSOL DEODORIZER SANITIZER

### ACTIVE INGREDIENTS:

n-Alkyl (60% C <sub>14</sub> , 30% C <sub>16</sub> , 5% C <sub>12</sub> , 5% C <sub>18</sub> ) dimethyl benzyl ammonium chlorides	5%
n-Alkyl (68% C <sub>12</sub> , 32% C <sub>14</sub> ) dimethyl ethylbenzyl ammonium chlorides	5%
Ethyl Alcohol	2.5%
INERT INGREDIENTS	87.5%
	100.00%

RINSOL is a concentrate and should be diluted before using.

NET CONTENTS \_\_\_\_\_ EPA Reg. No. 1791-28  
EPA Est. No. 1791-WA-1

**DANGER**  
**KEEP OUT OF REACH OF CHILDREN**

See other precautionary and first aid statements on left side panel.



USE DIR

#### Household

For sanitizing and disinfecting floors, walls and  
Locker Rooms, Garbage Pails, Sink Tops, Corridors  
solution with mop or cloth. Sanitize with 1 ounce  
ounces RINSOL to 5 gallons of water. Kills Staphylococcus  
For heavily soiled or contaminated areas, a pre

#### Hospitals

For floors, walls and other hard surfaces. In  
Garbage Pails, Telephones and Restrooms, use RINSOL  
with cloth, mop, or sponge. At this level RINSOL  
PRD-10. For heavily soiled or contaminated are

#### Food Processing Equipment

For sanitization of previously cleaned food processing  
RINSOL per 4 gallons of water to provide 200 ppm  
POTABLE WATER RINSE is required.

#### Dairies

To sanitize dairy equipment such as tanks, line  
equipment thoroughly. Then apply sanitizing solution  
water (200 ppm). At this level, NO POTABLE WATER  
of local Health Board.

This product when used on environmental, in  
water is effective against Influenza A2, Herpes S

**NORTHCO** Since 1919

**ANT RINSOL DEODORIZER**  
**SANITIZER**

ACTIVE INGREDIENTS:

30% C <sub>14</sub> , 30% C <sub>16</sub> , 5% C <sub>12</sub> , 5% C <sub>18</sub> ) benzyl ammonium chlorides .....	5%
5% C <sub>12</sub> , 32% C <sub>14</sub> ) dimethyl zyl ammonium chlorides .....	5%
Triol .....	2.5%
INGREDIENTS .....	87.5%
	100.00%

RINSOL is a concentrate and should be diluted before using.

CONTENTS \_\_\_\_\_ EPA Reg. No. 1791-28  
EPA Est. No. 1791-WA-1

**DANGER**

**KEEP OUT OF REACH OF CHILDREN**

For further precautionary and first aid statements on left side panel.

**North Coast Chemical Co., Inc.**  
Seattle, Washington 98108

**USE DIRECTIONS:**

**Household**

For sanitizing and disinfecting floors, walls and inanimate hard surfaces: For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and Shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce RINSOL to 4 gallons of water. Disinfect with 3 ounces RINSOL to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

**Hospitals**

For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use RINSOL at 3½ ounces per 5 gallons of water. Apply with cloth, mop, or sponge. At this level RINSOL will be effective against Pseudomonas aeruginosa PRD-10. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

**Food Processing Equipment**

For sanitization of previously cleaned food processing equipment and food utensils dilute 1 ounce RINSOL per 4 gallons of water to provide 200 ppm of active quaternaries. At this level, NO POTABLE WATER RINSE is required.

**Dairies**

To sanitize dairy equipment such as tanks, lines, pails, and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce RINSOL to 4 gallons of water (200 ppm). At this level, NO POTABLE WATER RINSE is required. Follow recommendations of local Health Board.

This product when used on environmental, inanimate, hard surfaces at 3½ ounces per gallon of water is effective against Influenza A2, Herpes Simplex, Adenovirus type 2, and Vaccinia Viruses.