

This product emulsifies readily, forming a stable emulsion when added to water. Slight agitation will maintain a uniform dispersion.

COTTON INSECTS: In early season applications for thrips and fleahoppers apply 1 pint per acre in 1 to 5 gallons of water and repeat at weekly intervals as long as infestations persist.

In pre-square applications for boll weevil control apply 1 quart per acre in 1 to 5 g dlons of water in 2 or 3 applications at weekly intervals at chopping time or as soon as the first weevils appear in the field. Consult local agricultural authorities regarding timing and adaptability of early season control programs in your locality.

For a mid-season control program for boll weevil apply 1 quart per acre in 2 to 5 gallons of water starting when up to 10% of the squares are penctured and repeat at 5 to 7 day intervals until 3 applications have been made or as long as weevil infestation persists.

108 S. Toombs St.

TOXAPHENE EMULSIFIABLE CONCENTRATE

(Contains 8 lbs. Toxaphene)

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene,
Chlorine Content 67-69%)
Methylated Aromatic Petroleum Derivatives
75.0%
INERT INGREDIENTS:
47.

Total 100.0%

WARNING! MAY BE FATAL IF SWALLOWED! Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after using. To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material. Avoid contamination of feed and foodstuffs. Cotton which has received late applications of this insecticide should not be grazed by dairy animals and animals being finished for slaughter. FLAMMABLE MIXTURE: Keep away from heat or open flame.

NET CONTENTS: ___ GALLONS PARRAMORE & GRIFFIN

For late season caterpillar app sure adequate Repeat applica

In the event is

important troi of the sell concerning the whether in ac