# COTTON-PEANUT DUST



20-40 TOXAPHENE - SULPHUR

### **ANALYSIS**

### **ACTIVE INGREDIENTS:**

SULPHUR FINENESS: Not less than 93% passing No. 325 USS Sieve

#### DIRECTION:

COTTON. For the control of bollweevils, bollworms and fleahopper. Use at the rate of 10 to 15 pounds per acre, depending on size of plant. For further directions as to time of application and amount consult Federal or State Experiment Stations or apply in accordance with accepted local practice.

PEANUTS: To control fall armyworms, cutworms and thrips on peanuts, apply 15 lbs. per acre. NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

## WARNING: KEEP OUT OF REACH OF CHILDREN.

Harmful it swallowed Harmful vapor and dust may be absorbed through the skin, may cause irritation of skin and eyes. Avoid inhaling or getting into eyes. Avoid skin contact. Avoid contamination of food and feed products. Keep clothing free from residues. For eyes, flush with water and get prompt medical attention.

#### ANTIDOTE:

INTERNAL: Give one tablespoon of salt in a glass of water and repeat until vemit is clear. EXTERNAL: Wash with soap and water

This product is toxic to fish and wildlife.

this produce is toxic to itsii and wining

Keep out of any body of water

Bird- feeding on treated accas may be killed

Do not apply when weather condition favor drift from area treated.

Do not reuse cars bar Empty bacs should be buried or burned. If burned stay well away floming smoke.

or by annuals being fine hed for laughter Do not feed treated ferage to dairy animals being fine hed for laughter Do not feed treated ferage to dairy animals or animals being fine hed for slaughter.

PARRAMORE & GRIFFIN

VALDOSTA, GEORGIA

# COTTON-PEANUT DUST



in the second of the second of

#### **ANALYSIS**

## **ACTIVE INGREDIENTS:**

Toxaphene (Technical Chlorinated Camphene)
Chlorine Content 67% to 69%
Sulphur, not less than
INFRT INGRIDIENTS:

TOTAL

TOTAL

SUI PHUR FINEMESS: Not less than 93% passing No. 325 USS Sieve

DIRECTION:

COTTON: For the control of bollweevils, bollworms and fleahopper. Use at the rate of 10 to 15 pounds per acre, depending on size of plant. For further directions as to time of application and amount consult Federal or State Experiment Stations or apply in accordance with accepted local practice.

PEANUTS: To control tall armyworms, cutworms and thrips on peanuts, apply 15 lbs, per sere NOT FOR USF OR STOR AGE IN OR AROUND THE HOME.

TACC !

VALDOSIA OFFICE