



MANEB 8



ACCEPTED

ACTIVE INGREDIENT:

JUL 16 1975 Maneb (Manganese ethylene bisdithiocarbamate) 6.4%
 (Manganese equivalent as metallic 16.5%)

INERT INGREDIENTS: 93.6%

**UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 1812-38**

TOTAL 100.0%

~~General Information~~ MANEB is sometimes referred to as MANZATE. Maneb is particularly effective against many of the fungus diseases of certain vegetable.

IMPORTANT: Never allow Maneb to become wet during storage. This could lead to chemical changes which reduce effectiveness as a fungicide and create vapors which may be flammable. Keep container closed when not in use.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CAUTION: KEEP OUT OF REACH OF CHILDREN

May cause irritation of eyes, nose, throat, and skin. Harmful if swallowed Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Keep away from fire or sparks. In case of contact, flush with plenty of water; for eyes, get medical attention.

DIRECTIONS

Watermelons: Downy mildew, anthracnose, Cercospora leaf spot and gummy stem blight—Apply as soon as plants begin to run or when disease threatens. Use 20-30 lbs per acre at 7 day intervals. When conditions are severe, apply at 3-day intervals.

Cucumbers: Downy mildew, anthracnose; same application recommendations as for watermelons.

Cantaloupe and Squash: Downy mildew—Apply as soon as plants begin to run or when disease threatens. Use 20-30 lbs. per acre and repeat at 7-10 day intervals as needed.

Cabbage Alternaria Leaf spot and Downy mildew—Use up to 40 lbs. per acre. Plant beds and direct seeded fields—apply 7-10 days after planting, or earlier if disease is present. Repeat at 3-day intervals until plants are set in the field or until field-seeded plants are thinned. Fields—apply when the disease threatens; repeat weekly as needed. Shorten interval to 3-5 days under severe conditions. Do not apply within 7 days of harvest.

Do not reuse container. Destroy when empty. E.P.A. EST. NO. 1812-GA-1

Parramore & Griffin

E.P.A. Reg. No. 1812-38

Valdosta, Ga.