



PEANUT DUST

COPPER - SULPHUR

ACCEPTED

131 JUL 1975

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 1812-43

THIS PACKAGE CONTAINS

ACTIVE INGREDIENTS:

Copper Oxide (as metallic) ----- 3.4%

Sulphur ----- 80.0%

INERT INGREDIENTS ----- 16.6%

Total ----- 100.0%

Sulphur Fineness: Not less than 93% passing No. 325 USS Sieve

RECOMMENDATIONS

Peanut Dust is a fungicide dust for the control of Peanut Leaf Spot. Consult your State Extension Service for complete details on the best peanut dusting program for your local conditions. These are general suggestions: Apply dust about sixty days after planting, or as soon as the spots appear on the leaf. First Dust—apply 15 to 20 pounds per acre. Later Dusts—apply 20 to 30 pounds per acre. A minimum of three to four applications should be made at two week intervals.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid inhaling dust. Avoid skin contact. Use respirator and goggles. Wash with soap and water after exposure. Keep clothing free from residues. Avoid contamination of foodstuffs. Sulphur may burn foliage when temperature is high—Do not make sulphur applications at such times.

ANTIDOTE: Induce vomiting by tickling the throat with a feather or the fingers and give an emetic of strong mustard water, or if available, use Syrup of Ipecac, one to two tablespoonsful, then give lime water, milk and honey, white of egg and mucilaginous drinks in copious draughts. Call physician or take patient to a hospital immediately.

CAUTION: Forage crops treated with this dust should not be fed to dairy animals or to animals to be slaughtered. Do not use on peanuts to be harvested within 30 days.

PARRAMORE & GRIFFIN

E.P.A. Reg. No. 1812--43

VALDOSTA, GEORGIA

E.P.A. Est. No. 1812-GA 1



PEANUT DUST

COPPER - SULPHUR

ACCEPTED

31 JUL 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 1812-13

THIS PACKAGE CONTAINS

ACTIVE INGREDIENTS:

Copper Oxide (as metallic) 3.4

Sulphur 80.0

INERT INGREDIENTS 16.6

Total 100.0

Sulphur Fineness: Not less than 93% passing No. 325 USS Sieve

RECOMMENDATIONS

Peanut Dust is a fungicide dust for the control of Peanut Leaf Spot. Consult your State Extension Service for complete details on the best peanut dusting program for your local conditions. There are general suggestions: Apply dust about 8 to 10 days after planting, or as soon as the spots appear on the leaf. First Dust - apply 15 to 20 pounds per acre. Later Dusts - apply 20 to 30 pounds per acre. A minimum of three to four applications should be made at two week intervals.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid inhaling dust. Avoid skin contact. Use respirator and goggles. Wash with soap and water after exposure. Keep clothing free from residues. Avoid contamination of foodstuffs. Sulphur may burn foliage when temperature is high. Do not make sulphur applications at such times.

ANTIDOTE: Induce vomiting by tickling the throat with a feather or the fingers and give an emetic of strong mustard water or if available, use Syrup of Ipecac, one to two tablespoonsful, then give lime water, milk and honey, white of egg and mucilaginous drinks in copious draughts. Call physician or take patient to a hospital immediately.

CAUTION: Forage crops treated with this dust should not be fed to dairy animals or to animals to be slaughtered. Do not use on peanuts to be harvested within 30 days.

PARRAMORE & GRIFFIN

Reg. No. 1812-13

VALDOSTA, GEORGIA