

ACCEPTED
5-28-71
1812-59

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene) (Chlorine Content 67% to 69%)	10.0%
Sulphur	70.0%
INERT INGREDIENTS:	20.0%

Apply This Product Only as Specified on This Label. **TOTAL** 100.0%

Sulphur Fineness: Not less than 93% Passing No. 325 USS Sieve

DIRECTIONS: PEANUTS—Sulphene is useful in the control of thrips, velvetbean caterpillars (peanutworm), cutworms, mites and red spiders.

Apply at the rate of 15 lbs per acre for early season control and up to 30 lbs. per acre later in the season. Make first application just as soon as the peanuts are up and REPEAT at weekly intervals. Do not apply within two weeks of expected harvest. **FIELDS WHERE PEANUTS HAVE BEEN GROWN IN EITHER OF THE TWO PRECEDING YEARS** will probably have a much earlier infection. It may be necessary to dust such fields two or three additional times.

CAUTION: Harmful if swallowed. Harmful vapor and dust may be absorbed through the skin causing irritation of skin and eyes. Avoid inhaling or getting into eyes. Avoid skin contact. Avoid contamination of food and feed products. Keep clothing free from residues. Use respirator and goggles. Wash with soap and water after exposure.

CAUTION: KEEP OUT OF REACH OF CHILDREN Do not feed treated forage to dairy animals or animals being finished for slaughter. Paper containers should be buried or burned. Stay well away from smoke.

ANTIDOTE: INTERNAL: Give one table spoon of salt in a glass of water and repeat until vomit is clear.

EXTERNAL: Wash with soap and water. This product is toxic to fish and wildlife. Birds feeding on treated area may be killed. Do Not apply where runoff is likely to occur. Keep out of any body of water. Do Not contaminate water by cleaning of equipment, or disposal of wastes.

Do not apply when weather conditions favor drift of dust from areas treated.

PARRAMORE & GRIFFIN

U.S.D.A. Reg. No. 1812-59

Valdosta, Georgia